

RECEIVED: March 24, 2019

REVISED: April 28, 2019

ACCEPTED: May 14, 2019

PUBLISHED: May 27, 2019

## $\mathcal{PT}$ deformation of Calogero-Sutherland models

Francisco Correa<sup>a</sup> and Olaf Lechtenfeld<sup>b</sup>

<sup>a</sup>*Instituto de Ciencias Físicas y Matemáticas, Universidad Austral de Chile,  
Casilla 567, Valdivia, Chile*

<sup>b</sup>*Institut für Theoretische Physik and Riemann Center for Geometry and Physics,  
Leibniz Universität Hannover,  
Appelstraße 2, 30167 Hannover, Germany*

*E-mail:* [francisco.correa@uach.cl](mailto:francisco.correa@uach.cl),  
[olaf.lechtenfeld@itp.uni-hannover.de](mailto:olaf.lechtenfeld@itp.uni-hannover.de)

ABSTRACT: Calogero-Sutherland models of  $N$  identical particles on a circle are deformed away from hermiticity but retaining a  $\mathcal{PT}$  symmetry. The interaction potential gets completely regularized, which adds to the energy spectrum an infinite tower of previously non-normalizable states. For integral values of the coupling, extra degeneracy occurs and a nonlinear conserved supersymmetry charge enlarges the ring of Liouville charges. The integrability structure is maintained. We discuss the  $A_{N-1}$ -type models in general and work out details for the cases of  $A_2$  and  $G_2$ .

KEYWORDS: Global Symmetries, Integrable Field Theories, Field Theories in Lower Dimensions, Integrable Hierarchies

ARXIV EPRINT: [1903.06481](https://arxiv.org/abs/1903.06481)

---

Contents

<b>1</b>	<b>Introduction and summary</b>	<b>1</b>
<b>2</b>	<b><math>\mathcal{PT}</math> deformation of Calogero-Sutherland models</b>	<b>2</b>
<b>3</b>	<b>The energy spectrum</b>	<b>4</b>
<b>4</b>	<b>Conserved charges and intertwiners</b>	<b>5</b>
<b>5</b>	<b>Details of the <math>A_2</math> model</b>	<b>7</b>
<b>6</b>	<b>Details of the <math>G_2</math> model</b>	<b>12</b>
<b>A</b>	<b>Potential-free frame</b>	<b>16</b>
<b>B</b>	<b>Ring of Laurent polynomials</b>	<b>16</b>
<b>C</b>	<b>Wave functions for the <math>A_2</math> model</b>	<b>17</b>
<b>D</b>	<b>Wave functions for the <math>G_2</math> model</b>	<b>19</b>
<b>E</b>	<b>Coefficients of the conserved charge <math>J_6</math></b>	<b>20</b>

---

**1 Introduction and summary**

Recently, integrable systems have been subjected more intensely to non-hermitian deformations, as has been reviewed in [1]. Specifically,  $\mathcal{PT}$  deformations of rational Calogero models and their spherical reductions have been analyzed in some detail [2–6]. It was found that the mathematical structures and tools pertaining to integrability are compatible with  $\mathcal{PT}$  deformations, as long as the latter respects the Coxeter reflection symmetries of the models. A bonus of certain  $\mathcal{PT}$  deformations is the complete regularization of the coincident-particle singularities of Calogero models, which leads to an enhancement of the Hilbert space of physical states by previously non-normalizable wave functions. For integral values of the Calogero coupling(s), most of the new states (called ‘odd’) are energy-degenerate with old ones (called ‘even’), giving rise to a  $\mathbb{Z}_2$ -grading and a conserved intertwiner  $Q$  on top of the Liouville integrals of motion. This  $Q$  is often called a ‘nonlinear supersymmetry’ charge, and it enhances the symmetry of the superintegrable system to a  $\mathbb{Z}_2$ -graded one [7].

Here, we carry our analysis of  $\mathcal{PT}$  deformed spherically reduced (or angular) Calogero models [6] over to the trigonometric or Calogero-Sutherland case. These models were completely solved more than 20 years ago [8–10] and describe  $N$  interacting identical particles on a circle or, equivalently, one particle moving on an  $N$ -torus subject to a particular external potential. The latter has inverse-square singularities on the hyperplanes corresponding to incident-particle locations. We shall see that, also here, there exists a deformation which

renders the potential nonsingular and retains the integrable structure, adding an infinite tower of new states to the energy spectrum and allowing for a nonlinear supersymmetry-operator mapping between ‘new’ and ‘old’ states. Employing  $\mathcal{PT}$ -deformed Dunkl operators, we construct the deformed intertwiners (shift operators increasing the coupling by unity) and analyze their action on the deformed energy eigenstates. We then find a set of deformed Liouville charges which intertwine homogeneously (like the Hamiltonian), so that their eigenvalues are preserved by the shift. Details are worked out for the three-body cases based on the  $A_2$  and  $G_2$  groups.

Our analysis can straightforwardly be extended to any (higher-rank) Coxeter group, but explicit expressions quickly become rather lengthy. While Dunkl operators and Liouville charges are known for all models (and  $\mathcal{PT}$ -deformed effortlessly), we are not aware of a general classification of Weyl *anti*-invariant<sup>1</sup> polynomials, which will be needed to extend the ring of Liouville charges to include all intertwiners. Also interesting is the exploration of the hyperbolic models and, finally, the elliptic ones. We have employed the most simple  $\mathcal{PT}$  deformation compatible with the symmetries of the model, but there exist other options. The two we shortly discuss in the context of the  $A_2$  model do not fully regularize the potential, but there may be other ones more suitable. A classification will be most welcome.

The paper is organized as follows. After defining the  $A_{N-1}$  Calogero-Sutherland model and introducing a suitable  $\mathcal{PT}$  deformation in section 2, we describe the energy spectrum including the eigenstates in section 3. The following section 4 is devoted to the construction of deformed conserved charges and intertwiners with the help of Dunkl operators. Sections 5 and 6 work out the details for the  $A_2$  and  $G_2$  cases, respectively. Explicit low-lying wave functions are listed in appendices C and D.

## 2 $\mathcal{PT}$ deformation of Calogero-Sutherland models

The  $N$ -particle model is governed by a rank- $N$  Lie algebra  $\mathfrak{g}$ . Translation invariance implies that  $\mathfrak{g} = A_1 \oplus \mathfrak{g}^\perp$ , where the  $A_1$  part represents the center of mass. It may be decoupled, but we retain it for the time being.

For  $A$ -type models,  $\mathfrak{g}^\perp = A_{N-1}$ . They describe  $N$  identical particles on a circle of circumference  $L$ , mutually interacting via a repulsive inverse-square two-body potential. We label the  $L$ -periodic particle coordinates as

$$q_i \in \mathbb{R}/L\mathbb{Z} \quad \text{with} \quad i = 1, 2, \dots, N, \quad \text{i.e.} \quad q_i \simeq q_i + n_i L \quad \text{for} \quad n_i \in \mathbb{Z}, \quad (2.1)$$

but it proves more convenient to pass to multiplicative coordinates

$$x_i = e^{2i\pi q_i/L}, \quad (2.2)$$

with the useful relations

$$\frac{\partial}{\partial q_i} = \frac{2i\pi}{L} \frac{x_i \partial}{\partial x_i}, \quad \sin \frac{\pi}{L}(q_i - q_j) = \frac{1}{2i} \frac{x_i - x_j}{\sqrt{x_i x_j}}, \quad \cot \frac{\pi}{L}(q_i - q_j) = i \frac{x_i + x_j}{x_i - x_j}. \quad (2.3)$$

---

<sup>1</sup>Meaning antisymmetric under any Weyl reflection.

The  $A_{N-1}$  Calogero-Sutherland model is defined by the Hamiltonian

$$\begin{aligned}
 H(g) &= -\frac{1}{2} \sum_{i=1}^N \frac{\partial}{\partial q_i^2} + \frac{\pi^2}{L^2} \sum_{i<j} \frac{g(g-1)}{\sin^2 \frac{\pi}{L}(q_i - q_j)} \\
 &= \frac{\pi^2}{L^2} \left[ 2 \sum_{i=1}^N \left( \frac{x_i \partial}{\partial x_i} \right)^2 - 4g(g-1) \sum_{i<j} \frac{x_i x_j}{(x_i - x_j)^2} \right].
 \end{aligned} \tag{2.4}$$

We remark on the invariance under  $g \mapsto 1-g$ . Rather than an  $N$ -body problem on a circle, this system may also be interpreted as a single particle moving on an  $N$ -torus  $T^N$  and subject to a particular external potential. The latter's singularities on the walls of the Weyl alcove restrict the particle motion to a fundamental domain in the  $A_1 \oplus A_{N-1}$  weight space. For later use we introduce the shorthand notation

$$\partial_i = \frac{\partial}{\partial x_i} \quad \text{and} \quad x_{ij} = x_i - x_j \tag{2.5}$$

as well as the totally antisymmetric degree-zero homogeneous rational function

$$\Delta = \prod_{i<j} x_{ij} \prod_k x_k^{-\frac{1}{2}(N-1)}. \tag{2.6}$$

To establish  $\mathcal{PT}$  symmetry, it is necessary to identify two involutions, a unitary  $\mathcal{P}$  and an anti-unitary  $\mathcal{T}$ , such that the deformed Hamiltonian is invariant under their product. While for the latter we take the standard choice of complex conjugation, the former leaves various possibilities. In this paper we shall choose  $\mathcal{P}$  to be parity flip of all coordinates,<sup>2</sup> thus

$$\mathcal{P} : q_i \mapsto -q_i \quad \Leftrightarrow \quad x_i \mapsto x_i^{-1} \quad \text{and} \quad \mathcal{T} : i \mapsto -i \quad \Leftrightarrow \quad x_i \mapsto x_i^{-1}. \tag{2.7}$$

The Hamiltonian (2.4) is parity symmetric, so a  $\mathcal{PT}$ -symmetric way of deforming can be induced by a  $\mathcal{PT}$ -covariant complex coordinate change. The obvious option is

$$q_i \mapsto q_i^\epsilon = q_i + i\epsilon_i \quad \Leftrightarrow \quad x_i \mapsto x_i^\epsilon = x_i e^{-2\epsilon_i} \quad \text{with} \quad \epsilon_i > 0, \tag{2.8}$$

implying

$$\mathcal{P} : q_i^\epsilon \mapsto -q_i^{-\epsilon} \quad \Leftrightarrow \quad x_i^\epsilon \mapsto (x_i^{-\epsilon})^{-1} \quad \text{and} \quad \mathcal{T} : i \mapsto -i \quad \Leftrightarrow \quad x_i^\epsilon \mapsto (x_i^{-\epsilon})^{-1}. \tag{2.9}$$

Thus, the multiplicative coordinate  $x_i^\epsilon$  is  $\mathcal{PT}$  invariant for any value of  $\{\epsilon_i\}$ . If we do not want to deform the center-of-mass degree of freedom we must impose the restriction  $\sum_i \epsilon_i = 0$ .

This deformation generically removes the singularities in the potential because

$$\frac{x_i^\epsilon - x_j^\epsilon}{\sqrt{x_i^\epsilon x_j^\epsilon}} = \sqrt{\frac{x_i}{x_j}} e^{-\epsilon_{ij}} - \sqrt{\frac{x_j}{x_i}} e^{+\epsilon_{ij}} \quad \text{and} \quad \frac{x_i^\epsilon - x_j^\epsilon}{x_i^\epsilon + x_j^\epsilon} = \frac{x_i e^{-\epsilon_{ij}} - x_j e^{+\epsilon_{ij}}}{x_i e^{-\epsilon_{ij}} + x_j e^{+\epsilon_{ij}}} \tag{2.10}$$

---

<sup>2</sup>For  $N=3$ , we shall ponder on some other choices later on.

never vanish for  $\epsilon_{ij} \equiv \epsilon_i - \epsilon_j$  different from zero. Therefore, the deformed Hamiltonian

$$H^\epsilon(g) = \frac{\pi^2}{L^2} \left[ 2 \sum_{i=1}^N (x_i \partial_i)^2 - 4g(g-1) \sum_{i < j} \frac{x_i^\epsilon x_j^\epsilon}{(x_{ij}^\epsilon)^2} \right] \quad (2.11)$$

no longer restricts the particle motion to a single Weyl alcove but allows it to range over the entire  $T^N$ . This space still being compact, the energy spectrum will remain discrete. Only in the  $L \rightarrow \infty$  limit we recover the rational Calogero model with its continuous spectrum. In the following, we drop the superscript ‘ $\epsilon$ ’ but understand to have a generic deformation turned on with  $\epsilon_{ij} \neq 0$ .

### 3 The energy spectrum

So far, the  $\mathcal{PT}$  deformation (2.8) is fully compatible with the integrability of the  $A$ -type Calogero-Sutherland model. It merely hides in the substitution  $x_i \mapsto x_i^\epsilon$ . This remains true for the energy spectrum: the known energy levels are unchanged under the deformation, and the eigenstates are obtained from the undeformed ones simply by again deforming the coordinates. However, due to the disappearance of the singularities in the potential, previously non-normalizable eigenstates become physical, adding extra states to the spectrum!

One popular way to completely label the energy eigenstates is by an  $N$ -tuple

$$\vec{n} = (n_1, n_2, \dots, n_N) \quad \text{with} \quad n_1 \geq n_2 \geq \dots \geq n_N \geq 0 \quad (3.1)$$

of quasiparticle excitation numbers. After removing the center-of-mass energy by boosting to its rest system one obtains

$$E_{\vec{n}}(g) = \frac{\pi^2}{L^2} \left[ 2 \sum_k n_k^2 - \frac{2}{N} \left( \sum_k n_k \right)^2 + 2g \sum_k (N+1-2k) n_k + \frac{1}{6} N(N-1)(N+1) g^2 \right]. \quad (3.2)$$

Due to translation invariance, this expression is invariant under a common shift  $n_k \rightarrow n_k + c$ . In order to remove this redundancy, we put  $n_N = 0$ , so that the sums over  $k$  run from 1 to  $N-1$  only. The energy is bounded from below, with the ground-state value

$$E_0 \equiv E_{\vec{0}} = \frac{1}{6} N(N-1)(N+1) g^2 \frac{\pi^2}{L^2} \quad \text{for} \quad g \geq 0 \quad (3.3)$$

but a different lower bound (minimally zero) for  $g < 0$ .

To study the degeneracy, we rewrite (3.2) as a sum of squares,

$$E_{\vec{n}}(g) = \frac{\pi^2}{L^2} \sum_{k=1}^{N-1} \frac{2}{k(k+1)} \left[ n_1 + n_2 + n_3 + \dots + n_{k-1} - k n_k - \left( N - \frac{1}{2} k(k+1) \right) g \right]^2 \quad (3.4)$$

$$= \frac{\pi^2}{L^2} \sum_{k=1}^{N-1} [\lambda_k - \mu_k g]^2 \quad \text{with}$$

$$\lambda_k = \frac{n_1 + n_2 + n_3 + \dots + n_{k-1} - k n_k}{\sqrt{k(k+1)/2}} \quad \text{and} \quad \mu_k = \frac{N - k(k+1)/2}{\sqrt{k(k+1)/2}}. \quad (3.5)$$

Any collection  $\vec{n}$  of quantum numbers uniquely yields an element  $\vec{\lambda} = (\lambda_1, \lambda_2, \dots, \lambda_{N-1})$  in a particular Weyl chamber of the  $A_{N-1}$  weight space  $\Lambda_{N-1}$ , and the energy of the corresponding state is given by the radius-squared of a circle in  $\Lambda_{N-1}$  centered at  $\vec{\mu}g$ . Since  $\vec{\mu}$  lies outside the Weyl chamber in question, for positive  $g$  the minimal distance from the circle center to the physical states is given by  $|\vec{\mu}|$  and represents the nonzero ground-state energy  $E_0(g)$ . So the degeneracy of a given energy level may be found by counting the number of physical weight lattice points on the appropriate “energy shell”.

The eigenfunctions of the Hamiltonian are given by Jack polynomials, on which there exists an extensive literature. They are of the form

$$\langle x|\vec{n}\rangle_g \equiv \Psi_{\vec{n}}^{(g)}(x) = R_{\vec{n}}^{(g)}(x) \Delta^g \quad \text{with} \quad R_{\vec{n}}^{(g)}(x) = P_{\vec{n}}^{(g)}(x) \prod_k x_k^{-p}, \quad (3.6)$$

$$\text{where} \quad x = (x_1, x_2, \dots, x_N), \quad |\vec{n}| = n_1 + n_2 + \dots + n_{N-1}, \quad p = |\vec{n}|/N,$$

and  $P_{\vec{n}}^{(g)}$  is a homogeneous permutation-symmetric polynomial of degree  $|\vec{n}|$  in  $x$ . The rational function  $R_{\vec{n}}^{(g)}$  is homogeneous of degree zero, but in the center-of-mass frame we have  $R_{\vec{n}}^{(g)} = P_{\vec{n}}^{(g)}$ . The function  $R_{\vec{n}}^{(g)}$  is a linear combination of symmetric basis functions

$$Q_{\vec{m}}^+(x) = x_1^{m_1-p} x_2^{m_2-p} \dots x_N^{m_N-p} + \text{all permutations} \quad (3.7)$$

$$\text{with} \quad p = |\vec{m}|/N \quad \text{and} \quad |\vec{m}| = |\vec{n}| - \ell N \quad \text{for} \quad \ell = 0, 1, \dots$$

(remember we put  $m_N = 0$ ). The coefficients are rational functions of the coupling  $g$ . At  $g=0$  only the leading term remains, and  $P_{\vec{n}}^{(0)} \propto Q_{\vec{n}}^+$ . Hence, all functions are Laurent polynomials in the variables  $y_i = x_i^{1/N}$ . The structure will become clear from the examples below.

Before the  $\mathcal{PT}$  deformation,  $\Delta \propto \prod_{i<j} x_{ij}$  vanishes at coinciding coordinate values (the Weyl-alcove walls), which renders the wave functions (3.6) non-square-integrable when  $g < 0$ . Therefore, the physical spectrum is empty there. However, due to the  $g \leftrightarrow 1-g$  symmetry of the Hamiltonian, we should consider the two “mirror values” of  $g$  together to form a single Hilbert space  $\mathcal{H}_g$ . Then, for a given value of  $g > \frac{1}{2}$ , a generic deformation (with all  $\epsilon_{ij}$  nonzero) will abruptly add a second infinite set of energy eigenstates to the spectrum, given by replacing  $g$  with  $1-g$ . Their energies are given by  $E_{\vec{n}}(1-g)$  from (3.2) or (3.4) for a second set of quantum numbers  $\vec{n}$ . This produces a second “energy shell”, which may carry states all the way down to zero energy (if  $\vec{\mu}(1-g)$  is located in the physical Weyl chamber). For particular (typical integer) values of  $g$  the two shells may possess simultaneous states, leading to an enhancement of energy degeneracy. We shall illustrate these features in the examples below.

#### 4 Conserved charges and intertwiners

A key tool in the construction of the spectrum and conserved charges is the Dunkl operator<sup>3</sup>

$$D_i(g) = \frac{\partial}{\partial q_i} - g \sum_{j(\neq i)} \cot \frac{\pi}{L}(q_i - q_j) s_{ij} = \frac{i\pi}{L} \left[ 2x_i \partial_i - g \sum_{j(\neq i)} \frac{x_i + x_j}{x_i - x_j} s_{ij} \right], \quad (4.1)$$

---

<sup>3</sup>For convenience we restrict to  $A$ -type models in the section. Section 6 deals with a more general case.

where the reflection  $s_{ij}$  acts on its right by permuting labels  $i$  and  $j$ . It obeys a simple commutation relation,

$$[D_i(g), D_j(g)] = -g^2 \frac{\pi^2}{L^2} \sum_k (s_{ijk} - s_{jik}), \quad \text{where } s_{ijk} = s_{ij}s_{jk} \quad (4.2)$$

effects a cyclic permutation of the labels  $i, j, k$ .

The importance of the Dunkl operator is twofold. First, any permutation-invariant (in general: Weyl-invariant) polynomial of some degree  $k$  in  $\{D_i\}$  will, when restricted to totally symmetric functions, give rise to a Liouville charge  $C_k$ , i.e. a conserved quantity in involution. A simple basis of this ring is provided by the Newton sums,

$$I_k(g) = \text{res} \sum_i D_i(g)^k \quad \Rightarrow \quad [I_k(g), I_\ell(g)] = 0, \quad (4.3)$$

where ‘res’ denotes the restriction to totally symmetric functions, giving

$$I_k(g) H(g) = H(g) I_k(g). \quad (4.4)$$

The total momentum and the Hamiltonian itself are the prime examples,

$$I_1(g) = iP = \frac{2i\pi}{L} \sum_i x_i \partial_i \quad \text{and} \quad I_2(g) = -2(H(g) - E_0(g)). \quad (4.5)$$

In the center of mass,  $P = 0$  of course. Only the first  $N$  charges are functionally independent; any  $I_{k>N}$  can be expressed in terms of these. The  $I_k$  for  $3 \leq k \leq N$  may be employed to lift the degeneracy of the state labelling by energy alone.

Second, the symmetric restriction of any *anti*-invariant polynomial of some degree  $k$  in the Dunkl operators will yield an *intertwining* operator (or shift operator)  $M_k(g)$ , obeying

$$M_k(g) H(g) = H(-g) M_k(g) = H(g+1) M_k(g). \quad (4.6)$$

The simplest such intertwiner is

$$M_{\bar{k}}(g) = \text{res} \frac{1}{\bar{k}!} \sum_{\text{permutations } i < j} \prod (D_i(g) - D_j(g)) \quad \text{with } \bar{k} = \frac{1}{2}N(N-1), \quad (4.7)$$

where the sum is over all permutations of the  $\bar{k}$  factors in the product. Comparing (4.6) with (3.4) it can be inferred that the action of  $M_{\bar{k}}(g)$  on the states is

$$M_{\bar{k}}(g) |\vec{n}\rangle_g \propto |\vec{n}-\vec{\delta}\rangle_{g+1} \quad \text{with } \vec{\delta} = (N-1, N-2, \dots, 2, 1), \quad (4.8)$$

which will vanish if the target quantum numbers no longer respect the restriction in (3.1). The shift operator translates the energy shell by the vector  $\vec{\mu}$ . Its repeated action will eventually get the state  $|\vec{n}\rangle_g$  to the edge of the physical Weyl chamber. Therefore, any state gets mapped to zero after a certain number of shifts. The adjoint intertwiner  $M_{\bar{k}}^\dagger(g) = M_k(-g)$  has the opposite action,  $\vec{n} \mapsto \vec{n} + \vec{\delta}$  while  $g \mapsto g-1$ . Since  $M_k$  has a nonzero kernel,  $M_{\bar{k}}^\dagger$  is not surjective.

The Liouville charges  $I_\ell$  together with the intertwiners  $M_k$  form a larger algebra, which is of interest. Beyond the total momentum and the Hamiltonian, the higher conserved charges (4.3) do not intertwine homogeneously but mix when  $M_k$  is passed through them,

$$M_k(g) I_\ell(g) = \left[ I_\ell(g+1) + \sum_{m<\ell} c_{km}(g) I_m(g+1) \right] M_k(g), \quad (4.9)$$

with some coefficients  $c_{km}(g)$  polynomial in  $g$ . However, it may be possible to find another basis  $\{C_k(g)\}$  which intertwines nicely,

$$M_k(g) C_\ell(g) = C_\ell(g+1) M_k(g), \quad (4.10)$$

meaning that the shift effected by  $M_k$  will map simultaneous eigenstates of the whole set  $\{C_k\}$  to each other. The composition  $M_k^\dagger(g)M_k(g)$  is by construction an element of the Liouville ring and thus can be expressed in terms of the  $I_k(g)$  (or  $C_k(g)$ ).

When  $g \in \mathbb{N}$ , the energy levels (let us call them ‘even’) are degenerate with some at coupling  $1-g \in -\mathbb{N}_0$  (call those ‘odd’). In this case, there exists an extra, odd, conserved charge<sup>4</sup>

$$Q_k(g) = M_k(g-1)M_k(g-2) \cdots M_k(2-g)M_k(1-g) \quad (4.11)$$

mapping  $\mathcal{H}_g$  to itself after fusing the spectra at couplings  $g$  and  $1-g$ . We note that the action of  $Q_k$  is well defined only after applying the  $\mathcal{PT}$  deformation, since the undeformed spectrum is empty for negative  $g$  values. The hidden supersymmetry operator  $Q_k$  maps between ‘even’ and ‘odd’ states in the joined spectrum, which arise from the originally positive and negative  $g$  values, respectively. Its square is a polynomial in the (even) Liouville charges, as will be seen in an example in (6.27) below.

## 5 Details of the $A_2$ model

In this section we work out the details of the simplest nontrivial case, which describes three particles on a circle interacting according to the  $A_2$  structure. For simplicity we put  $L = \pi$  from now on; the dimensions can easily be reinstated. The Hamiltonian in the center-of-mass frame then reads (deformation superscript ‘ $\epsilon$ ’ suppressed)

$$H(g) = 2 \sum_{i=1}^3 (x_i \partial_i)^2 - 4g(g-1) \sum_{i<j} \frac{x_i x_j}{(x_i - x_j)^2} = -\frac{1}{2} I_2(g) + 4g^2. \quad (5.1)$$

The other two conserved charges are

$$\begin{aligned} \frac{1}{i} I_1(g) &= 2 \sum_i x_i \partial_i = P \quad \text{and} \\ \frac{i}{8} I_3(g) &= (x_1 \partial_1)^3 - \left( 3g(g-1) \left[ \frac{x_1 x_2}{(x_1 - x_2)^2} + \frac{x_1 x_3}{(x_1 - x_3)^2} \right] + 2g^2 \right) x_1 \partial_1 + \text{cyclic}, \end{aligned} \quad (5.2)$$

---

<sup>4</sup>Each  $M$  factor could even carry a different index  $k$  but the resulting  $Q$  charges presumably differ only by Liouville-charge factors.



but  $I_4$  is already dependent,

$$I_4 = \frac{4}{3}I_3I_1 + \frac{1}{2}I_2^2 - I_2I_1^2 + \frac{1}{6}I_1^4 - g^2I_2 + \frac{1}{3}g^3I_1^2. \quad (5.3)$$

The energy formulæ (3.2) and (3.4) specialize to

$$\begin{aligned} E_{n_1, n_2}(g) &= \frac{4}{3}(n_1^2 + n_2^2 - n_1n_2) + 4g n_1 + 4g^2 \\ &= (n_1 + 2g)^2 + \frac{1}{3}(n_1 - 2n_2)^2 \quad \text{with } n_1 \geq n_2 \geq 0, \end{aligned} \quad (5.4)$$

and the ground state for  $g \geq 0$  is

$$\langle x|0, 0\rangle_g \equiv \Psi_0^{(g)}(x) = \Delta^g = \left( \frac{x_{12}x_{13}x_{23}}{x_1 x_2 x_3} \right)^g = (Q_{2,1}^-)^g \quad \text{with } E_0(g) = 4g^2. \quad (5.5)$$

Here, we introduced (for later purposes) the antisymmetric basis functions, so

$$Q_{m_1, m_2}^\pm = x_1^{m_1-p} x_2^{m_2-p} x_3^{-p} \pm x_1^{m_2-p} x_2^{m_1-p} x_3^{-p} + \text{cyclic} \quad \text{with } p = (m_1 + m_2)/3. \quad (5.6)$$

These Laurent polynomials (in  $x_i^{1/3}$ ) form a ring whose structure we detail in appendix B. In appendix C we list the explicit wave functions

$$\Psi_{n_1, n_2}^{(g)}(x) = R_{n_1, n_2}^{(g)}(x) \Psi_0^{(g)}(x) \quad (5.7)$$

(see (3.6)) for small values of  $n_1$ .

Each eigenstate  $|n_1, n_2\rangle$  corresponds to a point

$$(\lambda_1, \lambda_2) = \left( -n_1, \frac{1}{\sqrt{3}}(n_1 - 2n_2) \right) \quad \text{with } \lambda_1 \leq -\sqrt{3}|\lambda_2| \quad (5.8)$$

in a  $\frac{\pi}{3}$  wedge around the negative  $\lambda_1$  axis. The circles determining the energy eigenstates for couplings  $g$  and  $1-g$  are centered at

$$(2, 0)_g \quad \text{and} \quad (2, 0)_{(1-g)} \quad (5.9)$$


in  $\lambda$ -space, respectively. This is illustrated in figure 1. Since  $\mu_2 = 0$ , we have an obvious energy degeneracy for

$$|n_1, n_2\rangle_g \quad \text{and} \quad |n_1, n_1 - n_2\rangle_g \quad (5.10)$$


except for  $n_1 = 2n_2$ , of course. For  $g \geq 0$  there rarely appears higher degeneracy,<sup>5</sup> but at  $g < 0$  energy levels are up to 12-fold degenerate! This plethora of states becomes physical only after the  $\mathcal{PT}$  deformation and greatly enlarges the Hilbert space  $\mathcal{H}_g$  for any  $g > 0$ . Figure 2 displays the energy spectra with degeneracies for low levels and small integral values of  $g$ .

---

<sup>5</sup>Occasional energy values are triply or quadruply degenerate.



**Figure 1.** States (black dots) and “energy shells” (blue for  $g=3$ , red for  $g=-2$ ) in weight space for  $E = \frac{28}{3}$  (left) and  $E = \frac{148}{3}$  (right).



**Figure 2.** Low-lying energy spectrum and degeneracies (color-coded) for the  $A_2$  model at integer coupling  $g \in \mathbb{Z}$  in the range  $g \in [-4, +4]$ .

The basic intertwiner for the  $A_2$  model is of order three,

$$M_3(g) = \frac{1}{3} \text{res} (D_{12}D_{23}D_{31} + D_{23}D_{31}D_{12} + D_{31}D_{12}D_{23})(g) \quad (5.11)$$

with the obvious notation  $D_{ij} = D_i - D_j$ . It computes to

$$\begin{aligned} M_3(g) = & \partial_{12}\partial_{23}\partial_{31} - 2g \{ \cot q_{12}\partial_{23}\partial_{31} + \text{cyclic} \} + 4g^2 \{ \cot q_{12} \cot q_{23}\partial_{31} + \text{cyclic} \} \\ & - g(g-1) \{ \sin^{-2} q_{31}\partial_{31} + \text{cyclic} \} - 4g(2g-1) \{ \sin q_{12} \sin q_{23} \sin q_{31} \}^{-1} \\ & - 2g(3g^2 - g + 2) \cot q_{12} \cot q_{23} \cot q_{31} + 2g(g-1)(g+2) \{ \cot^3 q_{12} + \text{cyclic} \} \end{aligned} \quad (5.12)$$

where we abbreviated  $\partial_{ij} = \frac{\partial}{\partial q_i} - \frac{\partial}{\partial q_j}$  and  $q_{ij} = q_i - q_j$ . In terms of the multiplicative variables, the shift operator takes the form

$$\begin{aligned}
 M_3(g) \propto & (x_1 \partial_1 - x_2 \partial_2)(x_2 \partial_2 - x_3 \partial_3)(x_3 \partial_3 - x_1 \partial_1) \\
 & - g \left\{ \frac{x_1 + x_2}{x_1 - x_2} (x_2 \partial_2 - x_3 \partial_3)(x_3 \partial_3 - x_1 \partial_1) + \text{cyclic} \right\} \\
 & + g^2 \left\{ \frac{x_1 + x_2}{x_1 - x_2} \frac{x_2 + x_3}{x_2 - x_3} (x_3 \partial_3 - x_1 \partial_1) + \text{cyclic} \right\} - g(g-1) \left\{ \frac{x_3 x_1}{x_{31}^2} (x_3 \partial_3 - x_1 \partial_1) + \text{cyclic} \right\} \\
 & + 4g(2g-1) \frac{x_1 x_2 x_3}{x_{12} x_{23} x_{31}} - \frac{1}{4} g(3g^2 - g + 2) \frac{x_1 + x_2}{x_1 - x_2} \frac{x_2 + x_3}{x_2 - x_3} \frac{x_3 + x_1}{x_3 - x_1} \\
 & + \frac{1}{4} g(g-1)(g+2) \left\{ \left( \frac{x_1 + x_2}{x_1 - x_2} \right)^3 + \text{cyclic} \right\}. \tag{5.13}
 \end{aligned}$$

Its action on the states is

$$M_3(g) |n_1, n_2\rangle_g \propto |n_1 - 2, n_2 - 1\rangle_{g+1} \tag{5.14}$$

conserving the energy. In weight space it moves  $(\lambda_1, \lambda_2) \mapsto (\lambda_1 + 2, \lambda_2)$ .

For homogeneous intertwining relations, we redefine

$$C_1 = I_1, \quad C_2 = I_2 - 2E_0 = -2H \quad \text{and} \quad C_3 = I_3 - I_1 I_2 \tag{5.15}$$

which obey

$$M_3(g) C_k(g) = C_k(g+1) M_3(g). \tag{5.16}$$

This implies that the eigenvalue of  $C_3$  is also conserved under the shift action. Indeed, it is readily verified that

$$\begin{aligned}
 C_1 |n_1, n_2\rangle_g &= 0, \quad C_2(g) |n_1, n_2\rangle_g = -2 \left[ (n_1 + 2g)^2 + \frac{1}{3} (n_1 - 2n_2)^2 \right] |n_1, n_2\rangle_g, \\
 C_3(g) |n_1, n_2\rangle_g &= -\frac{8}{9} i (n_1 - 2n_2)(2n_1 - n_2 + 3g)(n_1 + n_2 + 3g) |n_1, n_2\rangle_g,
 \end{aligned} \tag{5.17}$$

which is compatible with the shift (5.14). The degeneracy reflection  $n_2 \mapsto n_1 - n_2$  flips the sign of  $C_3$ , so two such states can be discriminated by their  $C_3$  eigenvalues.

As expected, the composition of the intertwiner with its adjoint yields an expression in the Liouville charges,

$$\begin{aligned}
 M_3^\dagger M_3(g) \propto & 18I_3^2 - 36I_3 I_2 I_1 + 8I_3 I_1^3 - 3I_2^3 + 21I_2^2 I_1^2 - 9I_2 I_1^4 + I_1^6 + 2g^2(9I_2^2 - 6I_2 I_1^2 + I_1^4) \\
 & = 18C_3^2 + 8C_3 C_1^3 - 3C_2^3 + 3C_2^2 C_1^2 - C_2 C_1^4 + C_1^6 \\
 & \quad - 6g^2(3C_2 - C_1^2 + 8g^2)^2.
 \end{aligned} \tag{5.18}$$

Let us take a look at the extra degeneracy between even ( $g > 0$ ) and odd ( $g \leq 0$ ) states appearing when  $g \in \mathbb{Z}$ . The odd operator  $Q_3(g)$  mapping one to the other and defined in (4.11) is of order  $3(2g-1)$  and shifts the quantum numbers as

$$(n_1, n_2) \mapsto (n_1 - 4g + 2, n_2 - 2g + 1), \tag{5.19}$$

which produces a rather large kernel.  $Q_3$  commutes with all conserved charges  $I_k$ , so it keeps their eigenvalues. In weight space, it maps between the ‘even’ and ‘odd’ energy shells.

Finally, we briefly discuss two other kinds of  $\mathcal{PT}$  deformations in the  $A_2$ -model context, which we denote as ‘angular’ and ‘radial’, respectively. Different from the parity transformation in (2.7), which amounts to the outer conjugation automorphism of  $A_{N-1}$ , the angular and radial deformations are compatible with an elementary Coxeter reflection (or particle permutation), e.g.

$$\mathcal{P} : (q_1, q_2, q_3) \mapsto (q_2, q_1, q_3) \quad \text{and} \quad (x_1, x_2, x_3) \mapsto (x_2, x_1, x_3), \quad (5.20)$$

while  $\mathcal{T}$  remains complex conjugation.

The angular  $\mathcal{PT}$  deformation is homogeneous in the  $q_i$  coordinates, in contrast to the constant complex coordinate shift (2.8). It is induced by a complex orthogonal coordinate change,  $\vec{q} \mapsto \Gamma_\epsilon \vec{q}$  with  $\Gamma_\epsilon \in \text{SO}(3, \mathbb{C})$  modulo  $\text{SO}(3, \mathbb{R})$ , described in [6]. Explicitly,

$$q_i \mapsto q_i^\epsilon = \frac{1}{3} [(1+2 \cosh \epsilon) q_i + (1 - \cosh \epsilon - i\sqrt{3} \sinh \epsilon) q_j + (1 - \cosh \epsilon + i\sqrt{3} \sinh \epsilon) q_k] \quad (5.21)$$

with  $(i, j, k)$  being a cyclic permutation of  $(1, 2, 3)$ . This deformation does not entirely remove the singular loci of the potential given by

$$\begin{aligned} 0 = \sin q_{ij}^\epsilon &= \cosh(\sqrt{3} \sinh \epsilon q_k) \sin(\cosh \epsilon q_{ij}) + i \sinh(\sqrt{3} \sinh \epsilon q_k) \cos(\cosh \epsilon q_{ij}) \\ \Leftrightarrow \quad q_{ij} &= \frac{\ell \pi}{\cosh \epsilon} \wedge q_k = 0 \quad \text{for } \ell = 0, 1, 2, \dots, \end{aligned} \quad (5.22)$$

where again  $(i, j, k)$  are cyclic and we went to the center-of-mass frame, so  $q_{ik} + q_{jk} = -3q_k$ . For small enough  $\epsilon$ , only the origin  $\{q_i = 0\}$  remains singular, but with growing value of  $\epsilon$  extra singularities appear inside the Weyl alcove.

The radial  $\mathcal{PT}$  deformation is a nonlinear one,

$$q_i \mapsto q_i^\epsilon = q_i + i\epsilon q_{jk}/r \quad \text{with} \quad r^2 = q_{12}^2 + q_{23}^2 + q_{31}^2 \quad (5.23)$$

and  $(i, j, k)$  being cyclic once more. The remaining singularities occur at

$$q_{ij} = \ell \pi \wedge q_k = 0 \quad \text{for } \ell = 0, 1, 2, \dots, \quad (5.24)$$

and in addition one should average the potential,  $V \mapsto V_\epsilon + V_{-\epsilon}$  with  $V_\epsilon(q) = V(q^\epsilon)$ . Both cases can be parametrized jointly by writing

$$V_\epsilon(q_{ij}) = V(R(\epsilon) q_{ij} + iI(\epsilon) q_k) \quad \text{with} \quad \begin{cases} R(\epsilon) = \cosh \epsilon & \text{and} & I(\epsilon) = \sqrt{3} \sinh \epsilon \\ R(\epsilon) = 1 & \text{and} & I(\epsilon) = -3/r \end{cases} \quad (5.25)$$

for the angular and radial  $\mathcal{PT}$  deformation, respectively.

## 6 Details of the $G_2$ model

For a more complicated and non-simply-laced example, we turn to the  $G_2$  model [11, 12] for three particles on a circle and apply the constant-shift  $\mathcal{PT}$  deformation (2.8) but suppress it notationally. The  $G_2$  model adds to the previous two-body potential of the  $A_2$  case (5.1) a specific three-body interaction,

$$\begin{aligned}
 H(g) &= -\frac{1}{2} \sum_{i=1}^3 \frac{\partial}{\partial q_i^2} + \sum_{i<j} \frac{g_S(g_S-1)}{\sin^2(q_i-q_j)} + \sum_{i<j} \frac{3g_L(g_L-1)}{\sin^2(q_i+q_j-2q_k)} \\
 &= 2 \sum_{i=1}^3 (x_i \partial_i)^2 - 4g_S(g_S-1) \sum_{i<j} \frac{x_i x_j}{(x_i-x_j)^2} - 12g_L(g_L-1) \sum_{i<j} \frac{x_i x_j x_k^2}{(x_i x_j - x_k^2)^2},
 \end{aligned} \tag{6.1}$$

where the index ‘ $k$ ’ complements  $i$  and  $j$  to the triple  $(1,2,3)$ , there are two independent real couplings  $g_S$  and  $g_L$ , and we again put  $L = \pi$  for simplicity. The potential can be viewed as a sum of two copies of the  $A_2$  potential, with a relative coordinate rotation between them. The singular walls appear for

$$q_i - q_j = 0 \quad \text{and} \quad q_k = \frac{1}{3}(q_1+q_2+q_3) \quad \text{for } i, j, k \in \{1, 2, 3\}, \tag{6.2}$$

bounding the  $G_2$  Weyl chambers. The Weyl group is enhanced from  $S_3$  to  $D_6$ , generated by

$$s_{12} : (q_1, q_2, q_3) \mapsto (q_2, q_1, q_3) \quad \text{and} \quad j : (q_1, q_2, q_3) \mapsto \frac{2}{3}(q_1+q_2+q_3)(1, 1, 1) - (q_1, q_2, q_3) \tag{6.3}$$

and permutations, which for the  $x_i$  coordinates translates to

$$s_{12} : (x_1, x_2, x_3) \mapsto (x_2, x_1, x_3) \quad \text{and} \quad j : (x_1, x_2, x_3) \mapsto (x_1 x_2 x_3)^{2/3} \left( \frac{1}{x_1}, \frac{1}{x_2}, \frac{1}{x_3} \right). \tag{6.4}$$

The Hamiltonian (6.1) yields eigenvalues

$$\begin{aligned}
 E_{n_1, n_2}(g) &= \frac{4}{3}(n_1^2 + n_2^2 - n_1 n_2) + 4g_S n_1 + 4g_L(2n_1 - n_2) + 4(g_S^2 + 3g_L^2 + 3g_S g_L) \\
 &= (n_1 + 2g_S + 3g_L)^2 + \frac{1}{3}(n_1 - 2n_2 + 3g_L)^2 \quad \text{with } n_1 \geq 2n_2 \geq 0,
 \end{aligned} \tag{6.5}$$


and the ground-state wave function for  $g_S \geq 0$  and  $g_L \geq 0$  is

$$\langle x|0, 0\rangle_{g_S, g_L} \equiv \Psi_0^{(g_S, g_L)}(x) = \Delta_S^{g_S} \Delta_L^{g_L} \quad \text{with } E_0(g) = 4(g_S^2 + 3g_L^2 + 3g_S g_L), \tag{6.6}$$

where we introduced

$$\begin{aligned}
 \Delta_S &= \frac{(x_1-x_2)(x_1-x_3)(x_2-x_3)}{x_1 x_2 x_3} = Q_{2,1}^-, \\
 \Delta_L &= \frac{(x_1^2-x_2 x_3)(x_2^2-x_1 x_3)(x_3^2-x_1 x_2)}{x_1^2 x_2^2 x_3^2} = \frac{1}{2}(Q_{3,0}^+ - Q_{3,3}^+).
 \end{aligned} \tag{6.7}$$

In addition to the permutation symmetry inherited from the  $A_2$  model, we also have to impose (anti-)invariance under the additional (even) Coxeter element  $j$  in (6.4), which



**Figure 3.** States (black dots) and “energy shells” for  $(g_S, g_L) = (3, 2)$  (blue),  $= (-2, 2)$  (red),  $= (3, -1)$  (green) and  $= (-2, -1)$  (orange) for  $E = \frac{52}{3}$  (left) and  $E = 156$  (right).

implements an inversion in  $x$  space and flips the sign of the roots by a  $\pi$  rotation in the relative  $q$  space. Noting that  $j^2 = 1$  and  $[j, s_{ij}] = 0$  and

$$j : Q_{m_1, m_2}^\pm \mapsto \pm Q_{m_1, m_1 - m_2}^\pm \tag{6.8}$$

one sees that  $j$  flips the sign of both  $\Delta_S$  and  $\Delta_L$ , hence

$$j \Psi_0^{(g_S, g_L)}(x) = (-)^{g_S + g_L} \Psi_0^{(g_S, g_L)}(x). \tag{6.9}$$

We also deduce that the energy-degenerate states  $|n_1, n_2\rangle$  and  $|n_1, n_1 - n_2\rangle$  of the  $A_2$  model are related by the action of  $j$ . Therefore, only their sum or difference will be a  $G_2$ -model state, so the range of  $n_2$  can be restricted to  $n_2 \leq \frac{1}{2}n_1$ , as already claimed in (6.5).

The excited states then take the form

$$\Psi_{n_1, n_2}^{(g_S, g_L)}(x) = R_{n_1, n_2}^{(g_S, g_L)}(x) \Psi_0^{(g_S, g_L)}(x), \tag{6.10}$$

where the  $R_{n_1, n_2}^{(g_S, g_L)}$  are again particular Weyl-symmetric rational functions of degree zero. Appendix D contains a list of low-lying wave functions.

Each eigenstate  $|n_1, n_2\rangle$  corresponds to a point

$$(\lambda_1, \lambda_2) = \left( -n_1, \frac{1}{\sqrt{3}}(n_1 - 2n_2) \right) \quad \text{with} \quad \lambda_1 \leq -\sqrt{3}\lambda_2 \leq 0 \tag{6.11}$$

in a  $\frac{\pi}{6}$  wedge above the negative  $\lambda_1$  axis, in accord with one  $G_2$  Weyl chamber. The circles determining the energy eigenstates for couplings  $(g_S, g_L)$ ,  $(1 - g_S, g_L)$ ,  $(g_S, 1 - g_L)$  and  $(1 - g_S, 1 - g_L)$  are centered at

$$\begin{aligned} & (2g_S + 3g_L, -\sqrt{3}g_L), & (2 - 2g_S + 3g_L, -\sqrt{3}g_L), \\ & (3 + 2g_S - 3g_L, \sqrt{3}g_L - \sqrt{3}), & (5 - 2g_S - 3g_L, \sqrt{3}g_L - \sqrt{3}) \end{aligned} \tag{6.12}$$

in  $\lambda$ -space, respectively. This is illustrated in figure 3. After the  $\mathcal{PT}$  deformation, the Hilbert space  $\mathcal{H}_{g_S, g_L}$  comprises the four towers obtained from the four circles in figure 3. Again, for integral values of the couplings, the towers have matching energy levels, which greatly increases their degeneracy.

The  $G_2$  Dunkl operator is an extension of the  $A_2$  one (again with  $\{i, j, k\} = \{1, 2, 3\}$ ),

$$-iD_i(g) = 2x_i\partial_i - g_S \sum_{j(\neq i)} \frac{x_i+x_j}{x_i-x_j} s_{ij} - g_L \left( \sum_{j(\neq i)} \frac{x_i x_j + x_k^2}{x_i x_j - x_k^2} s_{ik} - 2 \frac{x_j x_k + x_i^2}{x_j x_k - x_i^2} s_{jk} \right) j. \quad (6.13)$$

The first two Newton sums in this Dunkl operator yield the conserved momentum and energy,

$$I_1 = \text{res} \sum_i D_i = iP \quad \text{and} \quad I_2 = \text{res} \sum_i D_i^2 = -2(H - E_0), \quad (6.14)$$

because  $\sum_i q_i$  and  $\sum_i q_i^2$  are not only permutation-symmetric but also invariant under the rotation  $j$  from (6.3). This, however, is not the case for  $\sum_i q_i^r$  when  $r \geq 3$ , but one can find a Weyl-invariant combination at order six,

$$\sigma_3 j \sigma_3 = \sigma_3 \left( \sigma_3 - 2\sigma_2\sigma_1 + \frac{4}{9}\sigma_1^3 \right) \quad \text{with} \quad \sigma_r = \sum_i q_i^r, \quad (6.15)$$

which generates another Liouville charge,

$$\begin{aligned} J_6 = & \text{res} \{ 5D_1^6 + 12D_1^5 D_2 + 12D_1^4 D_2^2 - 24D_1^4 D_2 D_3 + 10D_1^3 D_2^3 + 12D_1^3 D_2^2 D_3 \\ & + \text{permutations}(1,2,3) \}_{\text{symmetrized}} \\ = & 5(x_1\partial_1)^6 + 12(x_1\partial_1)^5 x_2\partial_2 + 12(x_1\partial_1)^4 (x_2\partial_2)^2 \\ & - 24(x_1\partial_1)^4 (x_2\partial_2)(x_3\partial_3) + 10(x_1\partial_1)^3 (x_2\partial_2)^3 + 12(x_1\partial_1)^3 (x_2\partial_2)^2 (x_3\partial_3) \\ & + c_{400} \cdot (x_1\partial_1)^4 + c_{310} \cdot (x_1\partial_1)^3 x_2\partial_2 + c_{220} \cdot (x_1\partial_1)^2 (x_2\partial_2)^2 + c_{211} \cdot (x_1\partial_1)^2 (x_2\partial_2)(x_3\partial_3) \\ & + c_{300} \cdot (x_1\partial_1)^3 + c_{210} \cdot (x_1\partial_1)^2 (x_2\partial_2) + c_{111} \cdot (x_1\partial_1)(x_2\partial_2)(x_3\partial_3) \\ & + c_{200} \cdot (x_1\partial_1)^2 + c_{110} \cdot (x_1\partial_1)(x_2\partial_2) + c_{100} \cdot (x_1\partial_1) + c_{000} + \text{permutations}(1,2,3) \end{aligned} \quad (6.16)$$

where symmetrization means Weyl ordering of every summand, and the coefficients  $c_{s_1 s_2 s_3}(x)$  are given in appendix E.

From (6.8) we see that the  $G_2$  model enjoys two separate intertwiners,

$$\begin{aligned} M_{3,S} &= \frac{1}{3} \text{res} (D_1 - D_2)(D_2 - D_3)(D_3 - D_1) + \text{cyclic}, \\ M_{3,L} &= \frac{1}{3} \text{res} (D_1 + D_2 - 2D_3)(D_2 + D_3 - 2D_1)(D_3 + D_1 - 2D_2) + \text{cyclic}, \end{aligned} \quad (6.17)$$

which independently shift by unity the couplings  $g_S$  and  $g_L$ , respectively,

$$\begin{aligned} M_{3,S} |n_1, n_2\rangle_{g_S, g_L} &\propto |n_1 - 2, n_2 - 1\rangle_{g_S + 1, g_L}, \\ M_{3,L} |n_1, n_2\rangle_{g_S, g_L} &\propto |n_1 - 3, n_2\rangle_{g_S, g_L + 1}. \end{aligned} \quad (6.18)$$

Their explicit form is

$$\begin{aligned}
M_{3,S} \propto & (x_1\partial_1 - x_2\partial_2)(x_2\partial_2 - x_3\partial_3)(x_3\partial_3 - x_1\partial_1) - g_S \left\{ \frac{x_1+x_2}{x_1-x_2} (x_2\partial_2 - x_3\partial_3)(x_3\partial_3 - x_1\partial_1) + \text{cyclic} \right\} \\
& + g_S^2 \left\{ \frac{x_1+x_2}{x_1-x_2} \frac{x_2+x_3}{x_2-x_3} (x_3\partial_3 - x_1\partial_1) + \text{cyclic} \right\} - g_S(g_S-1) \left\{ \frac{x_3x_1}{x_3^2} (x_3\partial_3 - x_1\partial_1) + \text{cyclic} \right\} \\
& + 9g_L(g_L-1) \left\{ \frac{x_2^2x_3x_1}{(x_2^2-x_3x_1)^2} (x_3\partial_3 - x_1\partial_1) + \text{cyclic} \right\} + 2g_S(4g_S-2+9g_L(g_L+1)) \frac{x_1x_2x_3}{x_{12}x_{23}x_{31}} \\
& - \frac{1}{4}g_S(3g_S^2-g_S+2) \frac{x_1+x_2}{x_1-x_2} \frac{x_2+x_3}{x_2-x_3} \frac{x_3+x_1}{x_3-x_1} + \frac{1}{4}g_S(g_S-1)(g_S+2) \left\{ \left( \frac{x_1+x_2}{x_1-x_2} \right)^3 + \text{cyclic} \right\} \\
& - 9g_Sg_L(g_L+1) \left\{ \frac{x_3+x_1}{x_3-x_1} \frac{x_2^2x_3x_1}{(x_2^2-x_3x_1)^2} + \text{cyclic} \right\}, \tag{6.19}
\end{aligned}$$

$$\begin{aligned}
M_{3,L} \propto & (x_1\partial_1 + x_2\partial_2 - 2x_3\partial_3)(x_2\partial_2 + x_3\partial_3 - 2x_1\partial_1)(x_3\partial_3 + x_1\partial_1 - 2x_2\partial_2) \\
& + 3g_L \left\{ \frac{x_2^2+x_3x_1}{x_2^2-x_3x_1} (x_1\partial_1 + x_2\partial_2 - 2x_3\partial_3)(x_2\partial_2 + x_3\partial_3 - 2x_1\partial_1) + \text{cyclic} \right\} \\
& + 9g_L \left\{ x_1x_2 \left[ \frac{(g_L-1)x_2x_3}{(x_2^2-x_1x_3)^2} + \frac{(g_L-1)x_1x_3}{(x_1^2-x_2x_3)^2} + \frac{2g_L(x_3^2+x_1x_2)}{(x_2^2-x_1x_3)(x_1^2-x_2x_3)} \right] (x_1\partial_1 + x_2\partial_2 - 2x_3\partial_3) + \text{cyclic} \right\} \\
& + 9g_S(g_S-1) \left\{ \frac{x_1x_2}{(x_1-x_2)^2} (x_1\partial_1 + x_2\partial_2 - 2x_3\partial_3) + \text{cyclic} \right\} - 27g_Lg_S(g_S-1) \left\{ \frac{x_1x_2}{(x_1-x_2)^2} + \text{cyclic} \right\} \\
& + 54g_L \left[ g_S(g_S-1) \frac{(x_1^2-x_2x_3+x_2^2-x_3x_1+x_3^2-x_1x_2)^3}{(x_1-x_2)^2(x_2-x_3)^2(x_3-x_1)^2} - g_L(g_L-3) \right] \frac{x_1^2x_2^2x_3^2}{(x_1^2-x_2x_3)(x_2^2-x_3x_1)(x_3^2-x_1x_2)} \\
& - 27g_L(g_L^2+g_L-2)x_1x_2x_3 \left\{ x_1 \frac{x_1^2+x_2x_3}{x_1^2-x_2x_3} + \text{cyclic} \right\} + 27g_L^3 \frac{(x_1^2+x_2x_3)(x_2^2+x_3x_1)(x_3^2+x_1x_2)}{(x_1^2-x_2x_3)(x_2^2-x_3x_1)(x_3^2-x_1x_2)}. \tag{6.20}
\end{aligned}$$

A better basis for the Liouville charges is

$$C_1 = I_1, \quad C_2 = I_2 - 2E_0 = -2H \quad \text{and} \tag{6.21}$$

$$\begin{aligned}
C_6 = & J_6 + 2(27g_L^2 + 24g_Lg_S + 8g_S^2)C_2C_1^2 - 9g_L^2C_2^2 - \frac{1}{9}(105g_L^2 + 96g_Lg_S + 32g_S^2)C_1^4 \\
& + 16(39g_L^4 + 72g_L^3g_S + 60g_L^2g_S^2 + 24g_Lg_S^3 + 4g_S^4)C_1^2 - 144g_L^4C_2 - 576g_L^6, \tag{6.22}
\end{aligned}$$

obeying homogeneous intertwining relations

$$\begin{aligned}
M_{3,S}(g_S, g_L) C_k(g_S, g_L) &= C_k(g_S+1, g_L) M_{3,S}, \\
M_{3,L}(g_S, g_L) C_k(g_S, g_L) &= C_k(g_S, g_L+1) M_{3,L}. \tag{6.23}
\end{aligned}$$

This is also signified by the action

$$C_6 |n_1, n_2\rangle_{g_S, g_L} = -\frac{64}{81} (3g_L+n_1-2n_2)^2 (6g_L+3g_S+2n_1-n_2)^2 (3g_L+3g_S+n_1+n_2)^2 |n_1, n_2\rangle_{g_S, g_L}. \tag{6.24}$$

The intertwining with their corresponding conjugates produces two polynomials in the Liouville charges,

$$\begin{aligned}
M_{3,S}^\dagger M_{3,S} &= -3C_6 - \frac{1}{6}C_1^6 + \frac{3}{2}C_2C_1^4 - \frac{7}{2}C_2^2C_1^2 + \frac{1}{2}C_2^3 + g_S^2(C_1^2 - 3C_2 - 8g_S^2)^2, \\
M_{3,L}^\dagger M_{3,L} &= 81C_6 + C_1^2(2C_1^2 - 9C_2)^2 + 81g_L^2(C_1^2 - 3C_2 - 24g_L^2)^2. \tag{6.25}
\end{aligned}$$



The intertwining operators also enable odd conserved charges when the couplings take integer values, in the form of the chain of operators

$$\begin{aligned} Q_{3,S}(g_S, g_L) &= M_{3,S}(g_S-1, g_L) M_{3,S}(g_S-2, g_L) \cdots M_{3,S}(2-g_S, g_L) M_{3,S}(1-g_S, g_L), \\ Q_{3,L}(g_S, g_L) &= M_{3,L}(g_S, g_L-1) M_{3,L}(g_S, g_L-2) \cdots M_{3,L}(g_S, 2-g_L) M_{3,L}(g_S, 1-g_L), \end{aligned} \quad (6.26)$$

which in the simplest non-trivial cases squares to the form of the polynomials in (6.25),

$$\begin{aligned} Q_{3,S}^2(2, g_L) &= \left( 3C_6 + \frac{1}{6}C_1^6 - \frac{3}{2}C_2C_1^4 + \frac{7}{2}C_2^2C_1^2 - \frac{1}{2}C_2^3 \right)^3 + \text{lower terms}, \\ Q_{3,L}^2(g_S, 2) &= \left( 81C_6 + 4C_1^2(C_1^2 - \frac{9}{2}C_2^2) \right)^3 + \text{lower terms}. \end{aligned} \quad (6.27)$$

## Acknowledgments

This work was partially supported by the Alexander von Humboldt Foundation, Fondecyt grant 1171475 and by the Deutsche Forschungsgemeinschaft under grant LE 838/12. This article is based upon work from COST Action MP1405 QSPACE, supported by COST (European Cooperation in Science and Technology). O.L. is grateful for the warm hospitality at CECs and Universidad Austral de Chile, where the main part of this work was done. F.C. is also grateful for the warm hospitality at Leibniz Universität Hannover.

## A Potential-free frame

We display some relations with the potential-free frame for the  $A_{N-1}$  model. By conjugating the Hamiltonian one can trade the potential for a first-order derivative term,

$$\begin{aligned} \tilde{H}(g) &= \Delta^{-g} H(g) \Delta^g = -\frac{1}{2} \sum_{i=1}^N \frac{\partial}{\partial q_i^2} - \frac{\pi}{L} g \sum_{i<j} \cot \frac{\pi}{L} (q_i - q_j) \left( \frac{\partial}{\partial q_i} - \frac{\partial}{\partial q_j} \right) \\ &= \frac{\pi^2}{L^2} \left[ 2 \sum_{i=1}^N \left( \frac{x_i \partial}{\partial x_i} \right)^2 + 2g \sum_{i<j} \frac{x_i + x_j}{x_i - x_j} (x_i \partial_i - x_j \partial_j) \right]. \end{aligned} \quad (A.1)$$

## B Ring of Laurent polynomials

For convenience we describe the multiplication rule for the Laurent polynomials

$$\begin{aligned} Q_{m_1, m_2}^\pm(y) &= y_1^{2m_1 - m_2} y_2^{2m_2 - m_1} y_3^{-m_1 - m_2} \pm y_1^{2m_2 - m_1} y_2^{2m_1 - m_2} y_3^{-m_1 - m_2} + \text{cyclic} \\ &= y_1^\alpha y_2^\beta y_3^\gamma + y_1^\beta y_2^\gamma y_3^\alpha \pm y_1^\gamma y_2^\alpha y_3^\beta \pm y_1^\alpha y_2^\gamma y_3^\beta \pm y_1^\beta y_2^\alpha y_3^\gamma \pm y_1^\gamma y_2^\beta y_3^\alpha \\ &=: [m_1, m_2]_\pm =: (\alpha, \beta, \gamma)_\pm \quad \text{with } \alpha + \beta + \gamma = 0. \end{aligned} \quad (B.1)$$

The relation between the parameters is

$$3m_1 = 2\alpha + \beta = -\beta - 2\gamma = \alpha - \gamma \quad \text{and} \quad 3m_2 = 2\beta + \alpha = -\alpha - 2\gamma = \beta - \gamma. \quad (B.2)$$

To remove the redundancy of the labelling, we stipulate that

$$m_1 \geq m_2 \geq 0 \quad \Leftrightarrow \quad \alpha \geq \beta \geq \gamma. \quad (\text{B.3})$$

It is immediate that  $[m_1, 0]_- = [m_1, m_1]_- = 0$ .

The obvious multiplication ( $\epsilon, \bar{\epsilon} \in \{+, -\}$ )

$$\begin{aligned} (\alpha, \beta, \gamma)_\epsilon \times (\bar{\alpha}, \bar{\beta}, \bar{\gamma})_{\bar{\epsilon}} &= (\alpha + \bar{\alpha}, \beta + \bar{\beta}, \gamma + \bar{\gamma})_{\epsilon\bar{\epsilon}} + (\alpha + \bar{\beta}, \beta + \bar{\gamma}, \gamma + \bar{\alpha})_{\epsilon\bar{\epsilon}} + (\alpha + \bar{\gamma}, \beta + \bar{\alpha}, \gamma + \bar{\beta})_{\epsilon\bar{\epsilon}} \\ &+ \bar{\epsilon}(\alpha + \bar{\alpha}, \beta + \bar{\gamma}, \gamma + \bar{\beta})_{\epsilon\bar{\epsilon}} + \bar{\epsilon}(\alpha + \bar{\beta}, \beta + \bar{\alpha}, \gamma + \bar{\gamma})_{\epsilon\bar{\epsilon}} + \bar{\epsilon}(\alpha + \bar{\gamma}, \beta + \bar{\beta}, \gamma + \bar{\alpha})_{\epsilon\bar{\epsilon}} \end{aligned} \quad (\text{B.4})$$

produces the law

$$\begin{aligned} [m_1, m_2]_\epsilon \times [n_1, n_2]_{\bar{\epsilon}} &= [m_1 + n_1, m_2 + n_2]_{\epsilon\bar{\epsilon}} + [m_1 - n_1 + n_2, m_2 - n_1]_{\epsilon\bar{\epsilon}} + [m_1 - n_2, m_2 + n_1 - n_2]_{\epsilon\bar{\epsilon}} \\ &+ \bar{\epsilon}[m_1 + n_1 - n_2, m_2 - n_2]_{\epsilon\bar{\epsilon}} + \bar{\epsilon}[m_1 + n_2, m_2 + n_1]_{\epsilon\bar{\epsilon}} + \bar{\epsilon}[m_1 - n_1, m_2 - n_1 + n_2]_{\epsilon\bar{\epsilon}}. \end{aligned} \quad (\text{B.5})$$

Even assuming  $m_1 \geq n_1$  (without loss of generality), the right-hand side may produce contributions  $[k_1, k_2]_\pm$  with  $k_1 < k_2$  or with  $k_2 < 0$ , which we outlawed. However, it is easy to see that

$$[k_1, k_2]_\pm = \pm[k_2, k_1]_\pm \quad \text{and} \quad [k_1, k_2]_\pm = \pm[k_1 - k_2, -k_2]_\pm, \quad (\text{B.6})$$

so one may employ the first relation in the first case and the second one in the second case to obtain an admissible result. Some examples are

$$\begin{aligned} [m_1, m_2]_\pm \times [0, 0]_+ &= 6[m_1, m_2]_\pm, \\ [m_1, m_2]_\pm \times [1, 0]_+ &= 2[m_1 + 1, m_2]_\pm + 2[m_1, m_2 + 1]_\pm + 2[m_1 - 1, m_2 - 1]_\pm, \\ [m_1, m_2]_\pm \times [1, 1]_+ &= 2[m_1 + 1, m_2 + 1]_\pm + 2[m_1, m_2 - 1]_\pm + 2[m_1 - 1, m_2]_\pm, \\ [m_1, m_2]_\pm \times [2, 0]_+ &= 2[m_1 + 2, m_2]_\pm + 2[m_1, m_2 + 2]_\pm + 2[m_1 - 2, m_2 - 2]_\pm, \\ [m_1, m_2]_\pm \times [2, 2]_+ &= 2[m_1 + 2, m_2 + 2]_\pm + 2[m_1, m_2 - 2]_\pm + 2[m_1 - 2, m_2]_\pm, \\ [m_1, m_2]_\pm \times [2, 1]_+ &= [m_1 + 2, m_2 + 1]_\pm + [m_1 + 1, m_2 + 2]_\pm + [m_1 + 1, m_2 - 1]_\pm \\ &+ [m_1 - 1, m_2 + 1]_\pm + [m_1 - 1, m_2 - 2]_\pm + [m_1 - 2, m_2 - 1]_\pm, \\ [m_1, m_2]_\pm \times [2, 1]_- &= [m_1 + 2, m_2 + 1]_\mp - [m_1 + 1, m_2 + 2]_\mp - [m_1 + 1, m_2 - 1]_\mp \\ &+ [m_1 - 1, m_2 + 1]_\mp + [m_1 - 1, m_2 - 2]_\mp - [m_1 - 2, m_2 - 1]_\mp. \end{aligned} \quad (\text{B.7})$$

## C Wave functions for the $A_2$ model

The  $A_2$  wave functions take the form

$$\langle x | n_1, n_2 \rangle_g \equiv \Psi_{n_1, n_2}^{(g)}(x) = R_{n_1, n_2}^{(g)}(x) \Delta^g = P_{n_1, n_2}^{(g)}(x) (x_1 x_2 x_3)^{-(n_1 + n_2)/3} \Delta^g, \quad (\text{C.1})$$

where  $x = (x_1, x_2, x_3)$  and  $P_{n_1, n_2}^{(g)}$  is a homogeneous permutation-symmetric Jack polynomial of degree  $n_1 + n_2$  in  $x$ . Passing to the more convenient variables

$$y_i = x_i^{1/3} \quad (\text{C.2})$$

the rational function  $R_{n_1, n_2}^{(g)}$  is a linear combination of symmetric basis functions

$$Q_{m_1, m_2}^+(y) = y_1^{2m_1 - m_2} y_2^{2m_2 - m_1} y_3^{-m_1 - m_2} + y_1^{2m_2 - m_1} y_2^{2m_1 - m_2} y_3^{-m_1 - m_2} + \text{cyclic} \quad (\text{C.3})$$

with  $m_1 + m_2 = n_1 + n_2 - 3\ell$  for  $\ell = 0, 1, 2, \dots$

The first few Laurent polynomials are

$$\begin{aligned} R_{0,0} &= Q_{0,0}^+ = 6, \\ R_{1,0} &= Q_{1,0}^+ = 2(y_1^2 y_2^{-1} y_3^{-1} + y_2^2 y_3^{-1} y_1^{-1} + y_3^2 y_1^{-1} y_2^{-1}), \\ R_{1,1} &= Q_{1,1}^+ = 2(y_1 y_2 y_3^{-2} + y_2 y_3 y_1^{-2} + y_3 y_1 y_2^{-2}), \\ R_{2,0} &= Q_{2,0}^+ + \frac{2g}{g+1} Q_{1,1}^+, \\ R_{2,1} &= Q_{2,1}^+ + \frac{g}{2g+1} Q_{0,0}^+, \\ R_{2,2} &= Q_{2,2}^+ + \frac{2g}{g+1} Q_{1,0}^+, \\ R_{3,0} &= Q_{3,0}^+ + \frac{6g}{g+2} Q_{2,1}^+ + \frac{2g^2}{(g+1)(g+2)} Q_{0,0}^+, \\ R_{3,1} &= Q_{3,1}^+ + \frac{g}{g+1} Q_{2,2}^+ + \frac{g(5g+3)}{2(g+1)^2} Q_{1,0}^+, \\ R_{3,2} &= Q_{3,2}^+ + \frac{g}{g+1} Q_{2,0}^+ + \frac{g(5g+3)}{2(g+1)^2} Q_{1,1}^+, \\ R_{3,3} &= Q_{3,3}^+ + \frac{6g}{g+2} Q_{2,1}^+ + \frac{2g^2}{(g+1)(g+2)} Q_{0,0}^+, \\ R_{4,0} &= Q_{4,0}^+ + \frac{8g}{g+3} Q_{3,1}^+ + \frac{6g(g+1)}{(g+2)(g+3)} Q_{2,2}^+ + \frac{12g^2}{(g+2)(g+3)} Q_{1,0}^+, \\ R_{4,1} &= Q_{4,1}^+ + \frac{3g}{g+2} Q_{3,2}^+ + \frac{g(7g+8)}{(g+2)(2g+3)} Q_{2,0}^+ + \frac{3g(4g+1)}{(g+2)(2g+3)} Q_{1,1}^+, \\ R_{4,2} &= Q_{4,2}^+ + \frac{g}{g+1} Q_{3,3}^+ + \frac{g}{g+1} Q_{3,0}^+ + \frac{4g(4g+1)}{(g+1)(2g+3)} Q_{2,1}^+ + \frac{g(10g^2+7g+3)}{2(g+1)^2(2g+3)} Q_{0,0}^+, \\ R_{4,3} &= Q_{4,3}^+ + \frac{3g}{g+2} Q_{3,1}^+ + \frac{g(7g+8)}{(g+2)(2g+3)} Q_{2,2}^+ + \frac{3g(4g+1)}{(g+2)(2g+3)} Q_{1,0}^+, \\ R_{4,4} &= Q_{4,4}^+ + \frac{8g}{g+3} Q_{3,2}^+ + \frac{6g(g+1)}{(g+2)(g+3)} Q_{2,0}^+ + \frac{12g^2}{(g+2)(g+3)} Q_{1,1}^+, \\ R_{5,0} &= Q_{5,0}^+ + \frac{10g}{g+4} Q_{4,1}^+ + \frac{20g(g+1)}{(g+3)(g+4)} Q_{3,2}^+ + \frac{20g^2}{(g+3)(g+4)} Q_{2,0}^+ + \frac{30g^2(g+1)}{(g+2)(g+3)(g+4)} Q_{1,1}^+, \\ R_{5,1} &= Q_{5,1}^+ + \frac{4g}{g+3} Q_{4,2}^+ + \frac{3g(g+1)}{(g+2)(g+3)} Q_{3,3}^+ + \frac{3g(3g+5)}{(g+2)(g+3)} Q_{3,0}^+ + \frac{2g(11g^2+20g+5)}{(g+2)^2(g+3)} Q_{2,1}^+ + \frac{6g^2(g+1)}{(g+2)^2(g+3)} Q_{0,0}^+. \end{aligned} \quad (\text{C.4})$$

From the paper of Lapointe and Vinet [8], one can see that the Jack polynomials can be also constructed in terms of modified Dunkl operators

$$\mathcal{D}_{i,\beta}(g) = x_i \left( \partial_i + g \sum_{i \neq j} \frac{1 - s_{ij}}{x_i - x_j} \right) + \beta. \quad (\text{C.5})$$

In terms of the combinations

$$\begin{aligned} B_1(g) &= x_1 \mathcal{D}_{1,g}(g) + x_2 \mathcal{D}_{2,g}(g) + x_3 \mathcal{D}_{3,g}(g), \\ B_2(g) &= x_1 x_2 \mathcal{D}_{1,g}(g) \mathcal{D}_{1,2g}(g) + x_2 x_3 \mathcal{D}_{2,g}(g) \mathcal{D}_{3,2g}(g) + x_3 x_1 \mathcal{D}_{3,g}(g) \mathcal{D}_{1,2g}(g) \end{aligned} \quad (\text{C.6})$$

the Jack polynomials are given by

$$P_{n_1, n_2}^{(g)} = B_2(g)^{n_2} B_1(g)^{n_1 - n_2} \cdot 1. \quad (C.7)$$

With the normalization

$$\Psi_{n_1, n_2}^{(g)} = \frac{1}{(g)_{n_1 - n_2} (g)_{n_2} (2g + n_1 - n_2)_{n_2}} P_{n_1, n_2}^{(g)} (x_1 x_2 x_3)^{-(n_1 + n_2)/3} \Delta^g \quad (C.8)$$

employing the Pochhammer symbol  $(a)_n$ , the action of the intertwiner takes the form

$$M_3(g) \Psi_{n_1, n_2}^{(g)} = n_2(n_1 + g)(n_1 - n_2) \Psi_{n_1 - 2, n_2 - 1}^{(g+1)}. \quad (C.9)$$

Clearly, the  $n_2=0$  and  $n_2=n_1$  states are annihilated by  $M_3(g)$ .

## D Wave functions for the $G_2$ model

In the variables  $y = (y_i) = (x_i^{1/3})$  the  $G_2$  wave functions take the form

$$\langle y | n_1, n_2 \rangle_{g_S, g_L} \equiv \Psi_{n_1, n_2}^{(g_S, g_L)}(y) = R_{n_1, n_2}^{(g_S, g_L)}(y) \Psi_0^{(g_S, g_L)}(y) \quad (D.1)$$

with  $\Psi_0^{(g_S, g_L)}$  given in (6.6) and (6.7) and  $R_{n_1, n_2}^{(g_S, g_L)}$  being a degree-zero rational function in  $y$ . It turns out that for  $n_1 + n_2$  not divisible by three  $\Psi$  does not in general factorize as  $R$  times  $\Psi_0$  in the ring of Laurent polynomials, so  $R$  is of a more general class. However, the full wave function  $\Psi$  can be expressed in terms of the symmetric and antisymmetric basis polynomials

$$Q_{m_1, m_2}^\pm(y) = y_1^{2m_1 - m_2} y_2^{2m_2 - m_1} y_3^{-m_1 - m_2} \pm y_1^{2m_2 - m_1} y_2^{2m_1 - m_2} y_3^{-m_1 - m_2} + \text{cyclic} \quad (D.2)$$

with  $m_1 + m_2 = n_1 + n_2 - 3\ell$  for  $\ell = 0, 1, \dots$ .

Some low-lying factorizable states are listed below. Beyond  $\Psi_{0,0}^{g_S, g_L} = Q_{0,0}^+ \Delta_S^{g_S} \Delta_L^{g_L}$  we have

$$\begin{aligned} \Psi_{1,0}^{g,0} &= [Q_{1,0}^+ + Q_{1,1}^+] \Delta_S^g, \\ \Psi_{2,0}^{g,0} &= \left( [Q_{2,0}^+ + Q_{2,2}^+] + \frac{2g}{g+1} [Q_{1,0}^+ + Q_{1,1}^+] \right) \Delta_S^g, \\ \Psi_{2,1}^{g,0} &= \left( Q_{2,1}^+ + \frac{g}{2g+1} Q_{0,0}^+ \right) \Delta_S^g, \\ \Psi_{3,0}^{g,0} &= \left( [Q_{3,0}^+ + Q_{3,3}^+] + \frac{12g}{g+2} Q_{2,1}^+ + \frac{4g^2}{(g+1)(g+2)} Q_{0,0}^+ \right) \Delta_S^g, \\ \Psi_{3,1}^{g,0} &= \left( [Q_{3,1}^+ + Q_{3,2}^+] + \frac{g}{g+1} [Q_{2,0}^+ + Q_{2,2}^+] + \frac{g(5g+3)}{2(g+1)^2} [Q_{1,0}^+ + Q_{1,1}^+] \right) \Delta_S^g, \\ \Psi_{4,0}^{g,0} &= \left( [Q_{4,0}^+ + Q_{4,4}^+] + \frac{8g}{g+3} [Q_{3,1}^+ + Q_{3,2}^+] + \frac{6g(g+1)}{(g+2)(g+3)} [Q_{2,0}^+ + Q_{2,2}^+] + \frac{12g^2}{(g+2)(g+3)} [Q_{1,0}^+ + Q_{1,1}^+] \right) \Delta_S^g, \\ \Psi_{4,1}^{g,0} &= \left( [Q_{4,1}^+ + Q_{4,3}^+] + \frac{3g}{g+2} [Q_{3,1}^+ + Q_{3,2}^+] + \frac{g(7g+8)}{(g+2)(2g+3)} [Q_{2,0}^+ + Q_{2,2}^+] + \frac{3g(4g+1)}{(g+2)(2g+3)} [Q_{1,0}^+ + Q_{1,1}^+] \right) \Delta_S^g, \\ \Psi_{4,2}^{g,0} &= \left( Q_{4,2}^+ + \frac{g}{g+1} [Q_{3,0}^+ + Q_{3,3}^+] + \frac{4g(4g+1)}{(g+1)(2g+3)} Q_{2,1}^+ + \frac{g(10g^2+7g+3)}{2(g+1)^2(2g+3)} Q_{0,0}^+ \right) \Delta_S^g, \end{aligned} \quad (D.3)$$

with  $\Delta_S = Q_{2,1}^-$  and  $\Delta_L = \frac{1}{2}(Q_{3,0}^+ - Q_{3,3}^+)$ .

In addition, one can infer that

$$\begin{aligned}
 \Psi_{2,1}^{0,g} &= Q_{2,1}^+ \Delta_L^g, \\
 \Psi_{3,0}^{0,g} &= \left( [Q_{3,0}^+ + Q_{3,3}^+] + \frac{2g}{2g+1} Q_{0,0}^+ \right) \Delta_L^g, \\
 \Psi_{4,2}^{0,g} &= \left( Q_{4,2}^+ + \frac{2g}{g+1} Q_{2,1}^+ \right) \Delta_L^g, \\
 \Psi_{5,1}^{0,g} &= \left( [Q_{5,1}^+ + Q_{5,4}^+] + \frac{2g}{g+1} Q_{4,2}^+ + \frac{g(5g+3)}{(g+1)^2} Q_{2,1}^+ \right) \Delta_L^g, \\
 \Psi_{6,0}^{0,g} &= \left( [Q_{6,0}^+ + Q_{6,6}^+] + \frac{4g}{g+1} Q_{6,3}^+ + \frac{4g(4g+1)}{(g+1)(2g+3)} [Q_{3,0}^+ + Q_{3,3}^+] + \frac{g(10g^2+7g+3)}{(g+1)^2(2g+3)} Q_{0,0}^+ \right) \Delta_L^g, \\
 \Psi_{6,3}^{0,g} &= \left( Q_{6,3}^+ + \frac{3g}{g+2} [Q_{3,0}^+ + Q_{3,3}^+] + \frac{2g^2}{(g+1)(g+2)} Q_{0,0}^+ \right) \Delta_L^g, \\
 \Psi_{7,2}^{0,g} &= \left( [Q_{7,2}^+ + Q_{7,5}^+] + \frac{3g}{g+2} [Q_{5,1}^+ + Q_{5,4}^+] + \frac{2g(7g+8)}{(g+2)(2g+3)} Q_{4,2}^+ + \frac{6g(4g+1)}{(g+2)(2g+3)} Q_{2,1}^+ \right) \Delta_L^g.
 \end{aligned} \tag{D.4}$$

For  $g_S, g_L \in \{0, 1\}$ , the model is free, so the states take a very simple form:

$$\begin{aligned}
 \Psi_{n_1, n_2}^{0,0} &= Q_{n_1, n_2}^+ + Q_{n_1, n_1 - n_2}^+, \\
 \Psi_{n_1, n_2}^{1,0} &= Q_{n_1+2, n_2+1}^- + Q_{n_1+2, n_1 - n_2 + 1}^-, \\
 \Psi_{n_1, n_2}^{0,1} &= Q_{n_1+3, n_2}^+ - Q_{n_1+3, n_1 - n_2 + 3}^+, \\
 \Psi_{n_1, n_2}^{1,1} &= Q_{n_1+5, n_2+1}^- - Q_{n_1+5, n_1 - n_2 + 4}^-.
 \end{aligned} \tag{D.5}$$

In the last two lines,  $\Delta_L$  can only be factored off in case  $n_1 + n_2$  is a multiple of three.

## E Coefficients of the conserved charge $J_6$

In order to express the coefficients  $c_{s_1 s_2 s_3}$  in (6.16) we introduce the combinations

$$\Omega_{ij} = \frac{x_i x_j}{(x_i - x_j)^2} \quad \text{and} \quad \Upsilon_k = \frac{x_k^2 x_i x_j}{(x_k^2 - x_i x_j)^2} \tag{E.1}$$

as well as

$$\tilde{\Omega}_{ij}^n = \frac{x_i + x_j}{x_i - x_j} \Omega_{ij}^n \quad \text{and} \quad \tilde{\Upsilon}_k^n = \frac{x_k^2 + x_i x_j}{x_k^2 - x_i x_j} \Upsilon_k^n, \tag{E.2}$$

where  $\{i, j, k\} = \{1, 2, 3\}$  and  $n$  is a positive integer. In this way the coefficients read

$$\begin{aligned}
 c_{400} &= -12 g_S (g_S - 1) (3\Omega_{23} + 4\Omega_{31}) - 36 g_L (g_L - 1) (2\Upsilon_1 + 4\Upsilon_3) - 75 g_L^2 - 84 g_L g_S - 28 g_S^2, \\
 c_{310} &= -12 g_S (g_S - 1) (7\Omega_{12} + 4\Omega_{23} + 9\Omega_{31}) - 36 g_L (g_L - 1) (8\Upsilon_1 + 5\Upsilon_2 + 11\Upsilon_3) \\
 &\quad - 4(69 g_L^2 + 60 g_L g_S + 20 g_S^2), \\
 c_{220} &= -12 g_S (g_S - 1) (3\Omega_{12} + 3\Omega_{31}) - 36 g_L (g_L - 1) (5\Upsilon_2 - 2\Upsilon_3) - 3(15 g_L^2 + 24 g_L g_S + 8 g_S^2), \\
 c_{211} &= -12 g_S (g_S - 1) (3\Omega_{23} - 9\Omega_{31}) + 36 g_L (g_L - 1) (5\Upsilon_1 + \Upsilon_3) + 24(3 g_L^2 + 3 g_L g_S + g_S^2),
 \end{aligned} \tag{E.3}$$

$$\begin{aligned}
 c_{300} &= -54g_S(g_S-1)\tilde{\Omega}_{31}+18g_L(g_L-1)(8\tilde{\Upsilon}_1-17\tilde{\Upsilon}_3), \\
 c_{210} &= 54g_S(g_S-1)(\tilde{\Omega}_{12}-4\tilde{\Omega}_{31})+18g_L(g_L-1)(48\tilde{\Upsilon}_1+6\tilde{\Upsilon}_2-27\tilde{\Upsilon}_3), \\
 c_{111} &= -18^2g_L(g_L-1)\tilde{\Upsilon}_3,
 \end{aligned} \tag{E.4}$$

$$\begin{aligned}
 c_{200} &= 36g_S(g_S-1)(2g_S+3g_L)^2\left(\frac{1}{2}\Omega_{23}+\Omega_{31}\right)+108g_L(g_L-1)(2g_S+3g_L)^2\left(\frac{1}{2}\Upsilon_1+\Upsilon_3\right) \\
 &\quad -54g_S(g_S-1)\Omega_{31}-36g_L(g_L-1)\left(2\Upsilon_1+\frac{47}{2}\Upsilon_3\right)+54g_S(g_S-1)(g_S+2)(g_S-3)\Omega_{31}^2 \\
 &\quad +36(g_L+1)g_L(g_L-1)(g_L-2)\left(6\Upsilon_1^2+\frac{51}{2}\Upsilon_3^2\right)-36\cdot 90g_L(g_L-1)\Upsilon_2^2 \\
 &\quad +54g_S^2(g_S-1)^2(4\Omega_{23}\Omega_{31}+\Omega_{31}\Omega_{12})+108g_S(g_S-1)g_L(g_L-1)(-4\Omega_{23}\Omega_{31}+\Omega_{31}\Omega_{12}) \\
 &\quad +54g_L^2(g_L-1)^2(-\Upsilon_2\Upsilon_3+16\Upsilon_3\Upsilon_1)+18(2g_S^4+12g_Lg_S^3+30g_L^2g_S^2+36g_L^3g_S+15g_L^4) \\
 &\quad +36g_S(g_S-1)g_L(g_L-1)(17\Omega_{12}\Upsilon_2+14\Omega_{23}\Upsilon_3+8\Omega_{31}\Upsilon_1+10\Upsilon_1\Omega_{23}+5\Upsilon_3\Omega_{12}), \tag{E.5}
 \end{aligned}$$

$$\begin{aligned}
 c_{110} &= 36g_S(g_S-1)(8g_S^2+24g_Sg_L+27g_L^2)\left(\frac{1}{2}\Omega_{12}+\Omega_{31}\right) \\
 &\quad +108g_L(g_L-1)(8g_S^2+24g_Sg_L+27g_L^2)\left(\Upsilon_2+\frac{1}{2}\Upsilon_3\right) \\
 &\quad -108g_S(g_S-1)\left(\frac{1}{2}\Omega_{12}+2\Omega_{31}\right)-36g_L(g_L-1)\left(50\Upsilon_2+\frac{11}{2}\Upsilon_3\right) \\
 &\quad +108g_S(g_S-1)(g_S+2)(g_S-3)\left(\frac{1}{2}\Omega_{12}^2+2\Omega_{31}^2\right) \\
 &\quad +108g_L^2(g_L-1)^2\left(14\Upsilon_2^2+\frac{5}{2}\Upsilon_3^2\right)-108g_L(g_L-1)(100\Upsilon_2^2+11\Upsilon_3^2) \\
 &\quad +54g_S^2(g_S-1)^2(\Omega_{23}\Omega_{31}+6\Omega_{31}\Omega_{12})+54g_S(g_S-1)g_L(g_L-1)(10\Omega_{23}\Omega_{31}-4\Omega_{31}\Omega_{12}) \\
 &\quad +54g_L^2(g_L-1)^2(19\Upsilon_1\Upsilon_2+74\Upsilon_3\Upsilon_1)+18(4g_S^4+24g_Lg_S^3+60g_L^2g_S^2+72g_L^3g_S+39g_L^4) \\
 &\quad +36g_S(g_S-1)g_L(g_L-1)(25\Omega_{12}\Upsilon_2+13\Omega_{23}\Upsilon_3+31\Omega_{31}\Upsilon_1+16\Upsilon_2\Omega_{31}+23\Upsilon_3\Omega_{12}), \tag{E.6}
 \end{aligned}$$

$$\begin{aligned}
 c_{100} &= -162g_S(g_S-1)g_L^2\tilde{\Omega}_{31}+54g_L(g_L-1)(9g_L^2-8)(\tilde{\Upsilon}_1-\tilde{\Upsilon}_3) \\
 &\quad +36\cdot 18(g_L(g_L-1)-8)g_L(g_L-1)(\tilde{\Upsilon}_1^2-\tilde{\Upsilon}_3^2) \\
 &\quad -162g_S(g_S-1)(g_S(g_S-1)-6g_L(g_L-1))(\tilde{\Omega}_{12}\Omega_{23}+\tilde{\Omega}_{31}\Omega_{12}) \\
 &\quad +162g_L^2(g_L-1)^2(8\tilde{\Upsilon}_1\Upsilon_2-13\tilde{\Upsilon}_2\Upsilon_3+5\tilde{\Upsilon}_3\Upsilon_1) \\
 &\quad -54g_S(g_S-1)g_L(g_L-1)(3\tilde{\Omega}_{12}\Upsilon_2-6\tilde{\Omega}_{23}\Upsilon_3+3\tilde{\Omega}_{31}\Upsilon_1+8\tilde{\Upsilon}_1\Omega_{23}-8\tilde{\Upsilon}_3\Omega_{12}) \\
 &\quad -54g_S(g_S-1)g_L(g_L-1)(\Omega_{12}\tilde{\Upsilon}_2+7\Omega_{23}\tilde{\Upsilon}_3-8\Omega_{31}\tilde{\Upsilon}_1-12\Upsilon_3\tilde{\Omega}_{12})
 \end{aligned} \tag{E.7}$$

$$\begin{aligned}
c_{000} = & 27g_L^6 + 81g_S(g_S-1)g_L^2(2g_L^2-1)\Omega_{12} + 81g_L(g_L-1)(6g_L^4-9g_L^2+4)(2g_L^2-1)\Upsilon_1 \\
& + 81g_L^2g_S(g_S-1)(g_S+2)(g_S-3)\Omega_{12}^2 + 81g_L(g_L-1)(g_L+2)(g_L-3)(9g_L^2-20)\Upsilon_1^2 \\
& + 324g_L(g_L-1)(g_L+2)(g_L-3)(g_L+4)(g_L-5)\Upsilon_1^3 \\
& + 162g_Lg_S^2(g_S-1)^2\Omega_{12}\Omega_{23} + 162g_L^2(g_L-1)^2(9g_L^2+9g_S(g_S-1)-8)\Upsilon_1\Upsilon_2 \\
& + 324g_Lg_S(g_S-1)(g_L-1)\Omega_{23} \left( \left( \frac{3}{2}g_L-1 \right) \Upsilon_1 + (3g_L^2-1)\Upsilon_3 \right) \\
& - 324g_Lg_S^2(g_S-1)^2(g_L-1)\Omega_{12}^2\Omega_{23} + 1944g_L^2(g_L-1)^2(g_L+2)(g_L-3)\Upsilon_1^2\Upsilon_2 \quad (E.8) \\
& + 324g_Lg_S(g_S-1)(g_L-1)\Omega_{12}^2(g_S(g_S-1)\Upsilon_1 + (g_S+2)(g_S-3)\Upsilon_3) \\
& + 648g_Lg_S(g_S-1)(g_L-1)(g_L+2)(g_L-3)\Upsilon_1^2(\Omega_{12}-\Omega_{23}) \\
& + 162g_L^2g_S(g_S-1)(g_L-1)^2\Omega_{12}\Upsilon_1(13\Upsilon_2+14\Upsilon_3) \\
& - 324g_Lg_S(g_S-1)(g_L-1)(2g_S(g_S-1)+3g_L(g_L-1)+8)\Omega_{12}\Omega_{23}\Upsilon_1 \\
& - 162g_Lg_S(g_S-1)(g_L-1)(3g_L(g_L-1)-16)\Omega_{12}\Omega_{31}\Upsilon_1 \\
& + 648g_L^2(g_L-1)^2(g_L^2-g_L+4)\Upsilon_1\Upsilon_2\Upsilon_3.
\end{aligned}$$

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

## References

- [1] A. Fring, *PT-symmetric deformations of integrable models*, *Phil. Trans. Roy. Soc. Lond. A* **371** (2013) 20120046 [[arXiv:1204.2291](https://arxiv.org/abs/1204.2291)] [[INSPIRE](#)].
- [2] A. Fring and M. Znojil, *PT-symmetric deformations of Calogero models*, *J. Phys. A* **41** (2008) 194010 [[arXiv:0802.0624](https://arxiv.org/abs/0802.0624)] [[INSPIRE](#)].
- [3] A. Fring and M. Smith, *Antilinear deformations of Coxeter groups, an application to Calogero models*, *J. Phys. A* **43** (2010) 325201 [[arXiv:1004.0916](https://arxiv.org/abs/1004.0916)] [[INSPIRE](#)].
- [4] A. Fring and M. Smith, *PT invariant complex  $E_8$  root spaces*, *Int. J. Theor. Phys.* **50** (2011) 974 [[arXiv:1010.2218](https://arxiv.org/abs/1010.2218)] [[INSPIRE](#)].
- [5] A. Fring and M. Smith, *Non-Hermitian multi-particle systems from complex root spaces*, *J. Phys. A* **45** (2012) 085203 [[arXiv:1108.1719](https://arxiv.org/abs/1108.1719)] [[INSPIRE](#)].
- [6] F. Correa and O. Lechtenfeld, *PT deformation of angular Calogero models*, *JHEP* **11** (2017) 122 [[arXiv:1705.05425](https://arxiv.org/abs/1705.05425)] [[INSPIRE](#)].
- [7] F. Correa, O. Lechtenfeld and M. Plyushchay, *Nonlinear supersymmetry in the quantum Calogero model*, *JHEP* **04** (2014) 151 [[arXiv:1312.5749](https://arxiv.org/abs/1312.5749)] [[INSPIRE](#)].
- [8] L. Lapointe and L. Vinet, *Exact operator solution of the Calogero-Sutherland model*, *Commun. Math. Phys.* **178** (1996) 425 [[q-alg/9509003](https://arxiv.org/abs/q-alg/9509003)] [[INSPIRE](#)].
- [9] A.M. Perelomov, É. Ragoucy and P. Zaugg, *Explicit solution of the quantum three-body Calogero-Sutherland model*, *J. Phys. A* **31** (1998) L559 [[hep-th/9805149](https://arxiv.org/abs/hep-th/9805149)] [[INSPIRE](#)].
- [10] W. García Fuertes, M. Lorente and A.M. Perelomov, *An elementary construction of lowering and raising operators for the trigonometric Calogero-Sutherland model*, *J. Phys. A* **34** (2001) 10963 [[math-ph/0110038](https://arxiv.org/abs/math-ph/0110038)].

- [11] C. Quesne, *Exchange operators and extended Heisenberg algebra for the three-body Calogero-Marchioro-Wolfes problem*, *Mod. Phys. Lett. A* **10** (1995) 1323 [[hep-th/9505071](#)] [[INSPIRE](#)].
- [12] C. Quesne, *Three-body generalization of the Sutherland model with internal degrees of freedom*, *Europhys. Lett.* **35** (1996) 407 [[hep-th/9607035](#)] [[INSPIRE](#)].