



Whole-Genome Sequences of Two New *Caballeronia* Strains Isolated from Cryoturbated Peat Circles of the Permafrost-Affected Eastern European Tundra

Stefanie A. Hetz,^a  Anja Poehlein,^b  Marcus A. Horn^a

^aSoil Microbiology, Institute of Microbiology, Leibniz University Hannover, Hannover, Germany

^bDepartment of Genomic and Applied Microbiology, Institute of Microbiology and Genetics, Georg August University of Göttingen, Göttingen, Germany

Stefanie A. Hetz and Anja Poehlein contributed equally to this work. Author order was determined alphabetically.

ABSTRACT Annotated genomes of *Caballeronia* strains SBC1 and SBC2 from acidic permafrost suggest a new species with a facultative lifestyle via oxygen and nitrate respiration. Thus, a contribution to nitrogen cycling in cold and low-pH environments is anticipated.

Cryoturbated peat circles (PCs) (62°57'E, 67°03'N) contain up to 2 mM pore water nitrate, emit large amounts of nitrous oxide (1, 2), and host new nitrate reducers (3). SBC1 and SBC2 were isolated from serial PC sediment (pH 4.2) dilutions by plating on semisolid modified R2A medium (1:10 diluted DSMZ 830 medium, 0.5% [wt/vol] K₂HPO₄, 7 g liter⁻¹ Gelrite [pH 6]) and incubating for 7 days at 15°C. Single colonies were picked and purified by restreaking four times onto the same medium.

High-molecular-weight DNA (HWD) for Nanopore sequencing was isolated with the MasterPure complete DNA and RNA purification kit (Biozym, Hessisch Oldendorf, Germany) from cells grown in liquid modified R2A medium (pH 5). HWD quality was checked on a Bioanalyzer 2100 using the DNA 12000 kit (Agilent Technologies, Waldbronn, Germany), and HWD was quantified with the Qubit double-stranded DNA (dsDNA) high-sensitivity (HS) assay kit (Life Technologies GmbH, Darmstadt, Germany); 1.5 μg HWD was used for library preparation employing the ligation sequencing kit 1D (SQK-LSK109) and the native barcode expansion kit (EXP-NBD114, barcode 15; Oxford Nanopore Technologies, Oxford, UK). Sequencing was performed for 72 h on the MinION Mk1B system with a SpotON flow cell R9.4.1 using MinKNOW v19.10.1, with Guppy v3.3.3 for base calling and demultiplexing. Totals of 67,997 reads with an average length of 4,873 bp (*N*₅₀, 27,443 bp) for SBC1 and 330,181 reads with an average length of 9,102 bp (*N*₅₀, 15,651 bp) for SBC2 were obtained.

Genomic DNA for Illumina shotgun sequencing was isolated via PCI extraction (4) and checked via spectrophotometry (DS-11; DeNovix, Inc., Wilmington, DE, USA). Illumina shotgun libraries were prepared using the Nextera XT DNA sample preparation kit, sequenced on a MiSeq system using reagent kit v3 with 600 cycles (2 × 300 bases; Illumina, San Diego, CA, USA), and resulted in totals of 3,249,515 (SBC1) and 2,281,633 (SBC2) paired-end reads per strain. Illumina reads were quality filtered using Trimmomatic v0.39 (5). Unicycler v0.4.6 (6) was used to perform a hybrid assembly, resulting in a closed circular chromosome (4,010,354 bp) and 4 closed plasmids (280,710 to 1,996,666 bp) for SBC1 and a closed circular chromosome (3,989,243 bp) and 7 closed plasmids (120,676 to 1,990,521 bp) for SBC2, as validated using Bandage v2.1 (7). Coverage was determined using Qualimap v2.2.1 (8) by mapping Illumina and Nanopore reads on the closed genomes using Bowtie 2 v2.3.5.1 (9) and minimap2 (10), respectively. Coverages for SBC1 and SBC2 were 98.4× and 62.4× (Illumina) and 62.8×

Citation Hetz SA, Poehlein A, Horn MA. 2020. Whole-genome sequences of two new *Caballeronia* strains isolated from cryoturbated peat circles of the permafrost-affected eastern European tundra. *Microbiol Resour Announc* 9:e00731-20. <https://doi.org/10.1128/MRA.00731-20>.

Editor J. Cameron Thrash, University of Southern California

Copyright © 2020 Hetz et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Marcus A. Horn, horn@ifmb.uni-hannover.de.

This is contribution 1 from the Institute of Microbiology, Leibniz University Hannover, to gain insights into hitherto unknown nitrate reducers from permafrost sediments.

Received 22 June 2020

Accepted 2 July 2020

Published 30 July 2020




FIG 1 Maximum composite likelihood tree of 16S rRNA genes aligned with MUSCLE (17) and rooted by midpoint rooting. Branches are scaled in terms of the expected number of substitutions per site. The percentage of trees in which the associated taxa clustered together is shown next to the branches. Initial trees for the heuristic search were obtained automatically by applying the neighbor-joining and BioNJ algorithms to a matrix of pairwise distances estimated using the maximum composite likelihood and the Tamura-Nei model (18) and then selecting the topology with a superior log likelihood value. The closest relative of SBC1 and SBC2 was *C. mineralivorans* from a fungal ectomycorrhizosphere in acidic and nutrient-poor forest soil (19). Tree construction was conducted with MEGA X, and GenBank accession numbers of 16S rRNA gene sequences are provided in parentheses next to species names (20).

and 162.8× (Nanopore), respectively. The overall GC contents (BioEdit v7.0.5.3 [11]) of SBC1 and SBC2 were 59.69% and 59.52%, respectively. Annotation with Prokka v1.14.0 (12) revealed the presence of 5 rRNA operons for both genomes, with 8,050 and 8,520 predicted protein-encoding genes and 59 and 60 tRNA genes for SBC1 and SBC2, respectively. Default parameters were used for all software.

SBC1 and SBC2 were affiliated with *Caballeronia mineralivorans* (Fig. 1); 53 to 57% of SBC1 and SBC2 genomes aligned with the *C. mineralivorans* genome, and the average nucleotide identity using the MUMmer algorithm (ANIm) (JSpeciesWS [13]) was 88.3%. The ANIm of SBC1 compared to SBC2 was 99.7%, suggesting that SBC1 and SBC2 represent a new species of the genus *Caballeronia* (14, 15). SBC1 and SBC2 encode multiple nitrate reductases of the *narG*, *napA*, and *nasA* types, as well as nitrite (*nirBD*) and nitric oxide (*norV*) reductases (Pathway Tools v23.0 [16]).

Data availability. These whole-genome shotgun projects have been deposited in DDBJ/ENA/GenBank under the accession numbers [CP049156.1](#) (chromosome) and [CP049157.1](#) to [CP049160.1](#) (plasmids) for SBC1 and under the accession numbers [CP049316.1](#) (chromosome) and [CP049317.1](#) to [CP049323.1](#) (plasmids) for SBC2. Bio-Project accession numbers for SBC1 and SBC2 are [PRJNA604524](#) and [PRJNA604525](#), and SRA accession numbers are [SRP250914](#) and [SRP250916](#), respectively.

ACKNOWLEDGMENTS

This work was financially supported by the Deutsche Forschungsgemeinschaft (grant DFG HO4020/3-1).

We are grateful to C. Biasi for providing sediment samples. We thank Sarah Teresa Schübler, Melanie Heinemann, and Natalie Röder for technical assistance.

REFERENCES

1. Repo ME, Susiluoto S, Lind SE, Jokinen S, Elsakov V, Biasi C, Virtanen T, Martikainen PJ. 2009. Large N₂O emissions from cryoturbated peat soil in tundra. *Nat Geosci* 2:189–192. <https://doi.org/10.1038/ngeo434>.
2. Marushchak ME, Pitkämäki A, Koponen H, Biasi C, Seppälä M, Martikainen PJ. 2011. Hot spots for nitrous oxide emissions found in different types of permafrost peatlands. *Glob Chang Biol* 17:2601–2614. <https://doi.org/10.1111/j.1365-2486.2011.02442.x>.
3. Palmer K, Biasi C, Horn MA. 2012. Contrasting denitrifier communities relate to contrasting N₂O emission patterns from acidic peat soils in arctic tundra. *ISME J* 6:1058–1077. <https://doi.org/10.1038/ismej.2011.172>.
4. Griffiths RI, Whiteley AS, O'Donnell AG, Bailey MJ. 2000. Rapid method for coextraction of DNA and RNA from natural environments for analysis of ribosomal DNA- and rRNA-based microbial community composition. *Appl Environ Microbiol* 66:5488–5491. <https://doi.org/10.1128/aem.66.12.5488-5491.2000>.
5. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
6. Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
7. Wick RR, Schultz MB, Zobel J, Holt KE. 2015. Bandage: interactive visualization of de novo genome assemblies. *Bioinformatics* 31:3350–3352. <https://doi.org/10.1093/bioinformatics/btv383>.
8. García-Alcalde F, Okonechnikov K, Carbonell J, Cruz LM, Götz S, Tarazona S, Dopazo J, Meyer TF, Conesa A. 2012. Qualimap: evaluating next-generation sequencing alignment data. *Bioinformatics* 28:2678–2679. <https://doi.org/10.1093/bioinformatics/bts503>.
9. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357–359. <https://doi.org/10.1038/nmeth.1923>.
10. Li H. 2018. minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34:3094–3100. <https://doi.org/10.1093/bioinformatics/bty191>.
11. Hall T. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* 41:95–98.
12. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
13. Richter M, Rosselló-Móra R, Oliver Glöckner F, Peplies J. 2016. JSpeciesWS: a Web server for prokaryotic species circumscription based on pairwise genome comparison. *Bioinformatics* 32:929–931. <https://doi.org/10.1093/bioinformatics/btv681>.
14. Richter M, Rosselló-Móra R. 2009. Shifting the genomic gold standard for the prokaryotic species definition. *Proc Natl Acad Sci U S A* 106:19126–19131. <https://doi.org/10.1073/pnas.0906412106>.
15. Varghese NJ, Mukherjee S, Ivanova N, Konstantinidis KT, Mavrommatis K, Kyrpides NC, Pati A. 2015. Microbial species delineation using whole genome sequences. *Nucleic Acids Res* 43:6761–6771. <https://doi.org/10.1093/nar/gkv657>.
16. Karp PD, Paley SM, Krummenacker M, Latendresse M, Dale JM, Lee TJ, Kaipa P, Gilham F, Spaulding A, Popescu L, Altman T, Paulsen I, Keseler IM, Caspi R. 2010. Pathway Tools version 13.0: integrated software for pathway/genome informatics and systems biology. *Brief Bioinform* 11:40–79. <https://doi.org/10.1093/bib/bbp043>.
17. Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32:1792–1797. <https://doi.org/10.1093/nar/gkh340>.
18. Tamura K, Nei M. 1993. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Mol Biol Evol* 10:512–526. <https://doi.org/10.1093/oxfordjournals.molbev.a040023>.
19. Uroz S, Oger P. 2017. *Caballeronia mineralivorans* sp. nov., isolated from oak-*Scleroderma citrinum* mycorrhizosphere. *Syst Appl Microbiol* 40:345–351. <https://doi.org/10.1016/j.syapm.2017.05.005>.
20. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol* 35:1547–1549. <https://doi.org/10.1093/molbev/msy096>.