



Cite this: *RSC Sustainability*, 2024, 2, 1058

Improving the stability and activity of a dye-decolourizing peroxidase using NADESs†

Maria Garbe, * Linnea Ute Lutz,  Leander Tom Lehmann, Theresa Strotmann, Ralf G. Berger and Franziska Ersoy

Natural Deep Eutectic Solvents (NADESs) are composed of naturally occurring compounds such as organic acids, amines and sugars; they are biodegradable and have low toxicities. NADESs have gained attention not only as environmentally friendly solvents, but also for their enzyme stabilization effects at high temperatures or osmotic stress. Six betaine-based NADESs were tested regarding their impact on enzymatic activity and thermal stability of a dye-decolourizing peroxidase from the basidiomycete *Pleurotus sapidus* (PsaPOX). Betaine : sucrose : water (2 : 1 : 10) and betaine : trehalose : water (4 : 1 : 14) had the best impact on the thermostability. Coloured silk was discoloured more efficiently by PsaPOX after addition of either NADES, which indicates a potential as stain-removal agents. The yield of the enzymatic transformation of *trans*-anethole to the aroma compound *p*-anisaldehyde increased by a factor of three with both tested NADESs. Overall, it was shown that betaine-based NADESs serve as sustainable media for reactions using PsaPOX and increase temperature stability as well as enzymatic activity. They might thus improve the industrial usability of PsaPOX.

Received 7th October 2023
Accepted 25th February 2024

DOI: 10.1039/d3su00359k

rsc.li/rscsus

Sustainability spotlight

The efficient use of resources, their substitution, reuse, or recycling play an increasingly important role in our society. Key are sustainable, biodegradable, low-toxicity, and naturally occurring substances for science and industry. Natural Deep Eutectic Solvents (NADES) possess these characteristics and are crucial for the green solvent technology. The present work demonstrates their utilisation for the temperature-stabilisation of a dye-decolourising peroxidase from the edible mushroom *Pleurotus sapidus*. Potential industrial applications like dye decolourisation and flavour production profited from the utilisation of NADES. NADES thus enable sustainable consumption and production, and supports measures to combat climate change, all of which are vital cornerstones of the UN's Sustainable Development Goals.

1. Introduction

Dye-decolourizing peroxidases (DyPs) (EC 1.11.1.19) represent a novel superfamily of heme peroxidases with the ability to oxidize a wide range of dyes and alkenes and thus numerous applications in various fields.^{1,2} One of the first described DyPs, from the basidiomycete *Thanatephorus cucumeris* (formerly *Geotrichum candidum*), decolourized 18 different reactive, acidic, and dispersing dyes, a majority of which were xenobiotics. The DyP exhibited higher decolourizing activities than classical peroxidases toward anthraquinone dyes.^{3–5} Some DyPs are also capable of degrading other natural- or azo dyes.^{6–9} As many dyes are toxic, mutagenic, and carcinogenic, and resistant to conventional wastewater treatments, DyPs are interesting tools for bioremediation.^{10–12} In addition, some DyPs have a redox potential which is high enough to enable the oxidation of lignin.¹³

Our group recently identified and characterized a DyP from *Pleurotus sapidus* (PsaPOX). It contained a non-canonical Mn²⁺-oxidation site on the protein surface and exhibited an alkene cleavage activity.⁷ The latter enabled its use as a biocatalyst to generate aromatic aldehydes with olfactory properties, such as veratraldehyde or *p*-anisaldehyde.^{7,14} These substances are used by the fragrance and flavor industries. It also bleached β -carotene and annatto, which is of interest for the detergent, milk, and baking industries.^{7,15,16} For industrial utilization, stabilizers are often necessary to improve the biocatalytic efficiency.^{17,18} One possibility is the use of natural deep eutectic solvents (NADESs).

NADESs are composed of two or more naturally occurring compounds. These can include sugars, amino acids, and organic acids, which, in combination, form a liquid phase at room temperature.^{19,20} Metabolic analyses have shown that NADESs naturally exist in organisms like plants: a system consisting of sucrose, fructose, and glucose protects cellular systems against freezing and drought.²¹ NADESs have several benefits over traditional organic solvents: they are biodegradable, have low toxicities, and are not easily flammable.²² In

Institute of Food Chemistry, Leibniz University Hannover, 30167 Hannover, Germany

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d3su00359k>



addition, they are simple to prepare and can be derived from renewable resources.²³

NADESS have been used for various applications, e.g., biomolecule extraction and separation, as environmentally friendly solvents for chemical reactions, and as protein stabilizers.^{24–27} They can be customized through an adjustment of their composition and properties to optimize conditions for different types of enzymes.^{28–33} Studies on horseradish peroxidase (HRP) showed that the NADES betaine : sorbitol : water (1 : 1 : 3) had the highest benefit because of its enzymatic activity. Alterations in the composition of the secondary structures illustrated the restructuring of the protein in the presence of NADESS through changes in the hydrogen bond network. Structural studies of HRP were conducted by determining the major secondary structure. The control contained 31% α -helices and 9% β -sheets, while, in the presence of NADESS, the α -helix content was extended by about 13%. This influenced the thermal stability of HRP positively.^{30,31} DESs and NADESS have shown advantageous effects on the activity and stability of various enzymes.²¹ Despite the limited industrial application of organophosphorus hydrolase due to its low thermal stability, promising results were observed by incubating the enzyme in ethaline and glyceline based-media.³⁴ Moreover, studies focusing on protein folding and thermostability have reported favourable outcomes. The presence of choline chloride–fructose improved the stability of the therapeutic protein human interferon- α 2 (IFN- α 2), showcasing the advantageous solvent environment provided by NADESS for storing therapeutic peptides.³⁵

In this work, we utilized betaine-based NADESS to improve the enzymatic activity and stability of PsaPOX. Six different NADESS were characterized regarding their effect on the relative enzymatic activity and the best two were chosen for further work. Their effect on temperature stability and different examples for potential industrial applications of PsaPOX were evaluated. Both the decolourization of silk and the biotransformation of *trans*-anethole to *p*-anisaldehyde were investigated. It is presumed that the use of NADESS will lead to an enhancement in the enzymatic activity of PsaPOX and influence its stability at high temperatures. These effects are expected to have an impact on the application possibilities.

2. Experimental

2.1. Materials

Chemicals were obtained from Carl-Roth (Karlsruhe, Germany), Sigma-Aldrich (Seelze, Germany) Merck (Darmstadt, Germany) and Oterra (Nienburg, Germany) in p.a. quality.

2.2. Synthesis of NADESS

The used NADESS were derived from Dai *et al.*, Khodaverdian *et al.*, and Tian *et al.*^{24,36,37} If NADESS were described as unstable in preliminary experiments, water was added to ensure long-term stability. NADESS were selected with substances known to have enzyme-stabilizing effects, such as sucrose and proline.^{21,38–40} All NADESS contained betaine as a hydrogen bond acceptor (HBA) and different combinations of hydrogen bond donors (HBD) (Table 1). The components were heated at 70 °C and 300 rpm until the solvents became stable and clear.

2.3. Protein expression and purification

The PsaPOX gene was expressed in *Komagataella phaffii* and the recombinant His-tagged enzyme was purified from the culture supernatant using Ni-NTA affinity chromatography as described before.^{7,41} The purity of the enzyme was determined using SDS (12% resolving, 4% stacking gel) and native PAGE.

2.4. Enzyme activity

The enzyme was prepared by mixing 22% NADES and 0.3 g per L peroxidase. The DyP was incubated at room temperature for 5 min. All assays were performed in triplicate. Blanks were carried out with water instead of the enzyme.

The peroxidase activity was determined photometrically (EON™ High Performance Microplate Spectrophotometer, Bio-Tek Instruments GmbH, Bad Friedrichshall, Germany) by monitoring the oxidation of 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid)diammonium salt (ABTS) as a substrate in the presence of hydrogen peroxide at 30 °C and 420 nm ($\epsilon_{420} = 3.6 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$) for 10 min. The samples were mixed with malonic acid buffer (50 mM, pH 4), 50 μM hydrogen peroxide and 0.5 mM ABTS in a total volume of 300 μL .

2.5. Thermostability

The thermostability of PsaPOX was tested in potassium phosphate buffer (50 mM) with 22% NADES. For the control, 22% 50 mM potassium phosphate buffer with the same pH value as the tested NADESS or 50 mM malonic acid buffer at pH 3.5, the pH-optimum of the enzyme, was added. PsaPOX was incubated at room temperature for 5 min at 200 rpm and 65, 75 or 80 °C for different intervals of time. For the stability tests at 65 °C, samples were taken after 2, 6, 10, 20, 30, 40 and 50 min, at 75 °C, samples were taken after 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55 and 60 min and at 80 °C after 2, 4, 6, 8 and 10 min. The samples were immediately placed on ice. All samples were brought to

Table 1 NADES compositions, including the molar ratios and abbreviations

Abbreviations	Component A	Component B	Component C	Molar ratio
BSucW	Betaine anhydrous	Sucrose	Water	2 : 1 : 10
BTrehW	Betaine anhydrous	Trehalose dihydrate	Water	4 : 1 : 14
BProGlu	Betaine anhydrous	D/L-Proline	Glucose	1 : 1 : 1
BSucPro	Betaine anhydrous	D/L-Proline	Sucrose	1 : 1 : 1
BMalPro	Betaine anhydrous	Malic acid	D/L-Proline	1 : 1 : 1
BMal	Betaine anhydrous	Malic acid	—	1 : 1



room temperature and their activity was determined using the ABTS assay as described above.

2.6. Decolourization assays

The decolourization of annatto, β -carotene, hispidin and laetiporic acid was evaluated. For β -carotene, a 9% (v/v) β -carotene emulsion was prepared with 1 g Tween 80 and 20 mL dichloromethane (DCM), and the solvent was removed by rotary evaporation.⁴²

9% (v/v) emulsion or annatto (Oterra, Nienburg), 100 μ M hydrogen peroxide, 25 mM manganese sulphate, 50 mM malonic acid buffer pH 4, 22% hispidin and laetiporic acid were extracted from fungal biomass using methanol as described elsewhere.^{43,44} 20 mL of these extracts were transferred into an aqueous emulsion as described for β -carotene.

2.7. Silk destaining

Coffee (Extra, Penny), red wine (Chianti Colli Senesi, 2020, Sviando, Italy), and annatto (Oterra, Nienburg) were used to dye 4.5–5.8 g silk. For the dyeing solution, 22.6 g soluble coffee powder was dissolved in 1.2 L boiling water. The coffee drink was cooled to room temperature. For the dyeing with red wine and annatto, 600 mL red wine or 4 mL annatto reagent was mixed with 200 or 800 mL water, respectively. 1 mL white vinegar was added to each dyeing solution. The silk was washed with water and heated with one of the dyeing solutions at 80 °C for 1 h. After cooling down to RT, the silk was washed for 2 min with water and dried for 24 h at RT. For the decolourization experiments, the silk was cut into 4 × 2 cm squares.

For decolourizing, a reaction mix containing 4 g NADES, 7.6 mL 50 mM malonate buffer pH 4, 1.58 mL 200 mM manganese sulphate, 4.82 mL purified enzyme (47 U L⁻¹), and 1.8 mL 1 mM hydrogen peroxide was prepared. Controls without the enzyme, as well as without NADESs were included. Three dyed silk pieces were covered with the reaction mix and incubated at RT for 30 min. After incubation, the silk pieces were washed for 2 min with deionized water and dried for 16 h at 30 °C. The colour intensity of the silk was measured after staining and destaining using a spectrophotometer (ColorLite sph900). All changes in colour intensity were calculated in comparison to changes of the controls.

2.8. Biotransformation of *trans*-anethole

The transformation of *trans*-anethole to *p*-anisaldehyde was carried out using 20 U L⁻¹ purified PsaPOX in 4 mL gas tight glass vials at a shaking rate of 200 rpm at 22 °C for 16 h in the absence of light. The reaction was performed in 50 mM Bis-Tris buffer pH 6, 100 μ M hydrogen peroxide, 25 mM manganese sulphate and 22% NADES or 50 mM potassium phosphate buffer with the same pH as the NADES sample. In a total volume of 1 mL, 1 μ L (6.7 mM) *trans*-anethole was added. Following the incubation period, *trans*-anethole and its resultant product, *p*-anisaldehyde, were extracted using 1 mL of hexane containing 100 mg per L (1 mM) cyclohexanol as an

internal standard (IS).⁷ Blanks were conducted without enzyme (chemical blank) or with heat-inactivated enzyme (2 hours at 95 °C). The organic phase was dried with anhydrous sodium sulphate and subsequently analyzed by gas chromatography (GC). GC measurements were conducted using a GC-2025 Shimadzu Instrument (Kyoto, Japan) equipped with a Cyclosil B column (30 m × 0.32 mm, 0.25 μ m, Agilent, Santa Clara, USA), a split/splitless injector port, and a flame ionization detection (FID) system. Hydrogen was employed as the carrier gas at a constant column flow rate of 2.5 mL per minute. A 1 μ L sample was injected *via* an autosampler and analyzed using the following method: 40 °C (3 minutes), a temperature ramp of 5 °C per minute until 220 °C, followed by a final hold time of 10 minutes. *Trans*-anethole and *p*-anisaldehyde were semi-quantified based on the area of the internal standard.

3. Results and discussion

3.1. Influence of NADESs on the enzymatic activity of PsaPOX

The influence of various NADESs on the activity of PsaPOX was determined. All NADESs were based on betaine as HBA, and sucrose, malic acid, glucose, proline and/or trehalose as HBD. PsaPOX was incubated in NADES-based media at 25 °C for 5 min and its relative activity was compared to that in the reference solution at the pH-optimum of 3.5 (100%, Fig. 1).⁷

PsaPOX exhibited the best activities in BSucW and BTrehW with 93.1 and 94.1%, respectively. Reduced activities were measured in BSucPro (59.2%), BMalPro (40.1%) and BMal (26.9%). No activity could be detected in BProGlu.

PsaPOX did not reach its maximal activity (100%) in any of the NADESs. This was due to the deviation from the pH optimum: Krahe *et al.* demonstrated the highest activity at pH




Fig. 1 Relative enzymatic activity of PsaPOX in six different NADESs using 50 mM malonic acid buffer (pH 3.5) as control (100%) at RT ($n = 3$). Statistical evaluation was carried out using unpaired *t*-tests in comparison to the control, * for $p < 0.05$, ** for $p < 0.01$, *** for $p < 0.001$ and **** for $p < 0.0001$, ns: not significant.



3.5, with more than 50% activity conserved between pH 3 and 5. At higher pH \leq 25% of the activity remained, with only 2.7% activity at pH 8.0, which corresponds to the pH of 7.9 of both BTrehW and BSucW.⁷ The shift in pH inevitably led to changes in the charge state and will have affected the protein structure. The addition of BSucW and BTrehW prevented this loss of activity. Instead, a high level of activity was maintained, indicating enzyme stabilization under significantly altered pH conditions. The results for BTrehW and BSucW confirm other studies which showed the ability of betaine-based NADESs to improve enzyme stability.^{24,31} Accordingly, BTrehW and BSucW were chosen for all subsequent analyses.

These two systems were tested with higher addition of water. This led to a change in viscosity and thus affected the enzyme activity. It can influence the formation of hydrogen bonds and may have contributed to an enhanced enzyme stabilization.³²

Due to the low pH optimum of PsaPOX, it was supposed that NADESs with a pH of 3–5 should further enhance the yields. Acids as HBD, such as lactic acid, should have reduced the pH value and thus enhanced the activity of PsaPOX, but did not show the desired effect (NADESs BMal and BMalPro) and were thus disregarded for further experiments.

Proline, which is known to protect enzymes in plants, also showed no positive effect on PsaPOX in this study (NADESs BSucPro and BMalPro).^{45,46}

Besides the water content of NADESs, their composition and the individual components also have an important influence on the relative activity. Altering the ratios changes the properties of the NADES and its interaction with the enzyme. For instance, certain DESs like ChCl–glycerol (2 : 1) or ChCl–ethylene glycol (2 : 1) had a negative impact on the relative activity of HRP but improved its thermal stability in higher concentrations.⁴⁷ The stability of NADESs themselves also needs to be ensured. NADESs containing sucrose and betaine (1 : 1) without water are not stable at room temperature.³⁶ It is crucial to find the correct concentrations of the respective NADES components. The composition can significantly influence their viscosity and stability, thereby impacting enzyme activity. For the tested NADESs and DyP in this study, the relative enzyme activity could not be increased, but the stability under different temperatures could be positively and significantly influenced.

For versatile peroxidases (VP) in ChCl–glycerol (1 : 2) and ChCl–urea (1 : 2), the enzyme activity decreased at different concentrations of DESs. At pH 6 and in the presence of 10% (v/v) ChCl–glycerol NADES, the highest enzyme activity was achieved with about twice that of the control.^{48,49} Similar effects could also be identified for the DyP.

3.2. Thermal stability of PsaPOX in NADESs

Both NADESs stabilized the enzyme activity of PsaPOX at 75 and 80 °C (Fig. 2). Without NADESs, a complete loss of enzyme activity occurred after 25 min at 75 °C. In contrast, samples containing BSucW and BTrehW showed measurable enzyme activities over the entire period of 60 min. After 15 minutes, activities of 32.1% (BSucW) and 37.5% (BTrehW) were measured in the NADESs, whereas only 1% remained in the

control. A similar trend was observed at 80 °C, where a loss of activity of up to 96% was detected within 4 min in the control sample, while the samples containing NADESs showed only 69.2 (BSucW) and 61.5% (BTrehW) loss. No significant changes were observed at 65 °C.

Many plants accumulate betaine and proline in stressful situations like heat or high salt conditions.^{50–53} It was shown that this protected enzymes from heat inactivation.³⁹ This was attributed to the formation of strong molecular interactions that potentially affect molecular mobility, which can also influence the thermal stability of proteins. The enzymatic activity of horseradish peroxidase at different temperatures was studied in betaine-based NADES.^{30,31} Several of them, including NADESs with trehalose, sucrose and water, resulted in a 60% increase in enzymatic activity. Betaine, trehalose, glycerol and water (2 : 1 : 3 : 5) kept HRP activity stable at 60 °C for 24 h.³⁰ One of the highest enzymatic activities was achieved in betaine–sorbitol–water (1 : 1 : 3), which achieved an about two-fold increase in enzyme activity.³¹ Also ChCl-containing DESs like ChCl–glycerol (2 : 1) and ChCl–ethylene glycol (2 : 1) were tested to improve thermal stability. Compared to the control, the NADESs slightly increased the relative activity of the HRP, compared to the control. Higher DES concentrations triggered the enzyme to be less active, but much more stable. For example, in the mixture of ChCl–urea (1 : 2), an enhancement in stability of up to 6.0-fold was measured.⁴⁷ For the tested NADESs in this study, thermal stability appeared to be positively influenced without any significant loss in relative activity. In a study by Wu *et al.*, a choline acetate-based NADES promoted both HRP activity and stability.⁴⁷

Not only did choline-based NADESs show effects on HRP, studies by Chang *et al.* showed a heat-stabilizing effect of sucrose on HRP.³⁸ In addition, trehalose is important for the induction of thermotolerance in yeast and other microorganisms.^{54,55} These positive effects of betaine, as well as trehalose and sucrose, were also observed here for PsaPOX.

Essential for the heat stability is the protein's secondary structure that undergoes restructuring in the presence of NADESs by establishing new hydrogen bond networks. CD (circular dichroism) spectroscopy has shown that the NADES which improved enzyme activity the most facilitated higher portions of α -helical structures.^{30,31} CD spectra showed that trypsin's secondary structure contained less α -helical area and a higher number of β -sheets folded into two domains in the NADES with betaine and trehalose.³⁷ Mamashli *et al.* measured ultraviolet CD spectra of versatile peroxidase with different DES contents at pH 4.5 and 7. The results showed that the α -helical content increased with the increase of DES concentration.⁴⁸ To explain the effect of NADESs on PsaPOX, the changes of the protein structure were determined in buffer with different concentrations of guanidine hydrochloride (GdnHCl) (Fig. S3†). The fluorescence of intrinsic aromatic amino acids increased more strongly in the control with higher concentrations of denaturing buffer. It has to be assumed that the addition of NADESs slows down the unfolding of PsaPOX. Gajardo-Parra *et al.* demonstrated that the addition of NADESs such as BSorbW resulted in an






Fig. 2 PsaPOX thermostability in two aqueous NADESs (BSucW and BTrehW) and in the reference buffer (control). Residual activity of the peroxidase was measured after incubation in phosphate buffer (50 mM, pH 8) in the absence (black) and presence of NADESs (22% BSucW (blue) or BTrehW (orange)) at (A) 65, (B) 75 or (C) 80 °C ($n = 3$). Statistical evaluation was carried out using unpaired t -tests in comparison to the respective control, * for $p < 0.05$, ** for $p < 0.01$, *** for $p < 0.001$ and **** for $p < 0.0001$, ns: not significant.

increase in the unfolding temperature of horseradish peroxidase (HRP). The findings suggest that interactions between NADESs and amino acids promote intermediate states during HRP folding, coupled with changes in the protein's secondary structure.³¹ It is assumed that these intermediate states are associated with a slower unfolding, which is also observed in DyPs.

Some NADESs also reduced enzyme flexibility, thus improving the active site stability. It is also known that NADESs generate an increase in water on the protein surface, which stabilizes the active site.³⁰ Which of these effects led to the increased stability of PsaPOX has to be addressed by future analyses, but the results are promising to run industrial processes over an extended period of time at elevated temperatures.




Fig. 3 Degradation of annatto, β -carotene, laetiporic acid, and hispidin with PsaPOX in the absence (grey) and presence of 22% NADESs BSucW (blue), BTrehW (orange) at 40 and 75 °C ($n = 3$). Statistical evaluation was carried out using unpaired t -tests in comparison to the control, * for $p < 0.05$, ** for $p < 0.01$, *** for $p < 0.001$ and **** for $p < 0.0001$, ns: not significant.






Fig. 4 Decolourising of silk with PsaPOX and 22% NADES or buffer. Due to the detaining effect of H_2O_2 , an additional control with hydrogen peroxide and without enzyme was added. (A) Decolourization after incubation at RT for 30 min using 47 U per L PsaPOX. (B) Decolourization of coffee-stained silk with higher enzymatic activities in the non-purified culture supernatant (100 U L^{-1}) and purified enzyme (47 U L^{-1}). (C) Decoloured silk (annatto) after 30 min at RT with 47 U per L PsaPOX. (D) Experimental setup with coloured silk (red wine). Statistical evaluation was carried out using unpaired *t*-tests in comparison to the control, * for $p < 0.05$, ** for $p < 0.01$, *** for $p < 0.001$ and **** for $p < 0.0001$, ns: not significant.

3.3. Decolourization of dyes in NADESs

PsaPOX belongs to a unique heme peroxidase family and is known for its ability to decolourize dyes.⁹ NADESs positively impacted the degradation of natural dyes such as β -carotene and annatto using PsaPOX (Fig. 3). Within 20 min at $40 \text{ }^\circ\text{C}$, approximately 32% β -carotene was degraded in NADESs, while only 24% degradation was observed in the buffer solution. The effect became even more pronounced at $75 \text{ }^\circ\text{C}$, where NADESs facilitated the degradation of 17% β -carotene, while only 4% degradation was evident in the control. For annatto, no degradation was measured in the control at $75 \text{ }^\circ\text{C}$, while 18% degradation occurred in both NADESs. Similar trends were observed with laetiporic acid. No significant differences between NADESs and control were observed for hispidin at $75 \text{ }^\circ\text{C}$ and at $40 \text{ }^\circ\text{C}$, a positive effect on dye degradation was shown only for BTrehW.

These findings further support the stabilizing effect of NADESs on enzymatic activities. Different studies have shown that the use of sugars as HBD leads to high viscosities due to the presence of a more pronounced intermolecular hydrogen-bond network. A decreased K_m was measured for a lipase in a NADES consisting of choline chloride and sucrose.⁵⁶ As the NADESs used in the present study contained sugars as HBD this is a possible explanation for the improved decolourisation.

In addition, the active site can be influenced by the change in hydrogen bonding patterns, leading to improved substrate

binding.²⁹ This can enhance substrate degradation and alter enzyme stability, as observed for PsaPOX. The literature about solvent–enzyme and NADES–substrate interactions is very limited. Nian *et al.* showed that the interaction of a lipase with NADESs increased the nucleophilic properties of the substrate, promoting the reaction.⁵⁷ A possible explanation is the effect of the increased water content in NADESs on solvated enzymes as described for lipases: the mobility of the enzyme increases with the hydration level of NADES.⁵⁸

The structural differences between hispidin as a polyketide and the other terpenoid dyes could have led to the formation of hydrogen bonds between NADESs and hispidin. A resultant substrate stabilization may prevent effective binding to the active site of PsaPOX or affect the reaction's activation energy and equilibrium. Comparable effects were discussed by Kovács *et al.*²⁹

In summary, it can be said that the improvement in decolourization may be a result of the enhanced stability of the DyP or an improved enzyme–substrate interaction facilitated by the NADES. It is evident that not all dyes are equally influenced by the interaction with the NADES and the DyP.

3.4. PsaPOX and NADESs as stain removers for textiles

Silk fabric was stained with red wine and coffee as examples for everyday stains and treated with a stain removal solution






Fig. 5 Alkene cleavage activity of PsaPOX on *trans*-anethole (6.7 mM) in the presence of 100 μ M H₂O₂ and 25 mM MnSO₄. Product concentrations after biotransformation to *p*-anisaldehyde ($n = 3$). Statistical evaluation was carried out using unpaired *t*-tests in comparison to the control, * for $p < 0.05$, ** for $p < 0.01$, *** for $p < 0.001$ and **** for $p < 0.0001$, ns: not significant.

(NADESS, PsaPOX, buffer, manganese sulphate, and hydrogen peroxide). In addition to the purified enzyme, the culture supernatant was tested as a stain remover, which would save the costs associated with enzyme purification for a potential industrial application. In almost all samples, a two-fold increase in textile decolourization was achieved by the addition of NADESS (Fig. 4A and B). Furthermore, a visually detectable improvement of the destaining of annatto treated silk was observed (Fig. 4C). The addition of hydrogen peroxide had a noticeable destaining effect; however, it was significantly lower than after the addition of PsaPOX. The culture supernatant with higher enzyme activities showed the activity-dependency of the decolourization (Fig. 4B). No inhibitory effect of the culture supernatant was detected, indicating that a combination with other destaining enzymes should be unproblematic. Purification of the enzyme is therefore not necessary, avoiding activity losses and reducing the cost of the approach.

3.5. Biotransformation of *trans*-anethole in NADESS

As shown above (Fig. 3), PsaPOX showed alkene cleavage activity towards the natural dyes β -carotene, annatto and laetiporic acid in NADESS. An improved cleavage was also observed for *trans*-anethole. The product concentration of *p*-anisaldehyde was increased by a factor of 3 with both NADESS (Fig. 5).

As demonstrated in previous experiments, enzyme stabilization may have occurred during the 16 hour incubation period with the NADESS. There was no significant difference between BTrehW and BSucW in this regard. Increased cleavage of *trans*-anethole increased the production of *p*-anisaldehyde, which can

be used for the synthesis of pharmaceuticals or as a fragrance in the food industry. Another explanation for the improved biotransformation in NADESS is the formation of hydrogen bonds for stabilizing the enzyme–substrate complex.²⁹ Similar results have already been achieved in cannabinoid production. Compared to aqueous systems, the use of NADESS resulted in a two-fold increase of cannabinoid production with a tetrahydrocannabinolic acid synthase and more than a three-fold increase with a cannabichromenic acid synthase.⁵⁹

A study by Yang *et al.* explored the feasibility of chemobiocatalytic production of furfuryl alcohol from D-xylose using tandem catalysis, the deep eutectic solvent betaine : benzene-sulfonic acid and recombinant *Escherichia coli* DCF cells. Application of 2.5 wt% DES BE : BA catalyzed D-xylose at 170 °C for 0.5 h, yielding 51.1% furfural.⁶⁰ NADESS hold promise as sustainable media for improving product yields in various industries.

4. Conclusions

This study established that several betaine-based NADESS enhanced PsaPOX stability at temperatures of 75 and 80 °C and a pH of 7.9. Moreover, NADESS improved the degradation of polyene dyes such as β -carotene and annatto and removal of coffee and red wine stains from silk samples. Alkene cleavage of *trans*-anethole to *p*-anisaldehyde was also enhanced.

The potential of PsaPOX as a biocatalyst for the degradation of β -carotene, as a stain remover and for the generation of aromatic aldehydes with olfactory properties was shown. For all tested applications, NADESS served as sustainable media with a stabilizing effect. Recently, molecular-scale modelling of NADESS and enzymes was established as an alternative to determine the optimal NADES for a specific enzyme based on the predicted intermolecular interactions.^{29,61} This should simplify and broaden the application of NADESS to improve enzymatic reactions.

Author contributions

M. G. – conceptualization, data curation, investigation, methodology, writing: original draft; L. U. L. – investigation; L. T. L. – investigation, data curation, formal analysis, T. S. – investigation, data curation, formal analysis; R. G. B. and F. E. – methodology, supervision, funding acquisition, writing – review and editing. All authors have read and agreed to the final version of the manuscript.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- 1 T. Caldas, N. Demont-Caulet, A. Ghazi and G. Richarme, *Microbiology*, 1999, **145**(pt 9), 2543–2548.
- 2 Y. Sugano, R. Muramatsu, A. Ichiyanagi, T. Sato and M. Shoda, *J. Biol. Chem.*, 2007, **282**, 36652–36658.



- 3 S.-J. Kim, K. Ishikawa, M. Hirai and M. Shoda, *J. Ferment. Bioeng.*, 1995, **79**, 601–607.
- 4 M. Shoda, *Commun. Agric. Appl. Biol. Sci.*, 2003, **68**, 269–274.
- 5 S. Kim and M. Shoda, *Appl. Environ. Microbiol.*, 1999, **65**, 1029–1035.
- 6 G. Rajhans, S. Sen, A. Barik and S. Raut, *J. Appl. Microbiol.*, 2020, **129**, 1633–1643.
- 7 N.-K. Krahe, R. G. Berger and F. Ersoy, *Molecules*, 2020, **25**, 69–85.
- 8 C. Lauber, T. Schwarz, Q. K. Nguyen, P. Lorenz, G. Lochnit and H. Zorn, *AMB Express*, 2017, **7**, 1–15.
- 9 D. I. Colpa, M. W. Fraaije and E. van Bloois, *J. Ind. Microbiol. Biotechnol.*, 2014, **41**, 1–7.
- 10 A. Bafana, S. S. Devi and T. Chakrabarti, *Environ. Rev.*, 2011, **19**, 350–371.
- 11 W.-Y. Li, F.-F. Chen and S.-L. Wang, *Protein Pept. Lett.*, 2010, **17**, 621–629.
- 12 A. K. Singh, M. Bilal, H. M. N. Iqbal, A. S. Meyer and A. Raj, *Sci. Total Environ.*, 2021, **777**, 145988.
- 13 C. Liers, C. Bobeth, M. Pecyna, R. Ullrich and M. Hofrichter, *Appl. Microbiol. Biotechnol.*, 2010, **85**, 1869–1879.
- 14 N.-K. Krahe, R. G. Berger, M. Witt, H. Zorn, A. B. Omarini and F. Ersoy, *Int. J. Mol. Sci.*, 2021, **22**(1363), 91–113.
- 15 R. T. Szweda, K. Schmidt and H. Zorn, *Eur. Food Res. Technol.*, 2013, **237**, 377–384.
- 16 C. Harrel, *Ind. Eng. Chem.*, 1952, **44**, 95–100.
- 17 A. S. Bommarius and M. F. Paye, *Chem. Soc. Rev.*, 2013, **42**, 6534–6565.
- 18 M. Pätzold, S. Siebenhaller, S. Kara, A. Liese, C. Sylдатk and D. Holtmann, *Trends Biotechnol.*, 2019, **37**, 943–959.
- 19 A. P. Abbott, G. Capper, D. L. Davies, R. K. Rasheed and V. Tambyrajah, *Chem. Commun.*, 2003, 70–71.
- 20 Y. Liu, J. B. Friesen, J. B. McAlpine, D. C. Lankin, S.-N. Chen and G. F. Pauli, *J. Nat. Prod.*, 2018, **81**, 679–690.
- 21 S. M. Taklimi, A. Divsalar, B. Ghalandari, X. Ding, M. L. Di Gioia, K. A. Omar and A. A. Saboury, *J. Mol. Liq.*, 2023, **377**, 121562.
- 22 A. P. Abbott, D. Boothby, G. Capper, D. L. Davies and R. K. Rasheed, *J. Am. Chem. Soc.*, 2004, **126**, 9142–9147.
- 23 A. Paiva, R. Craveiro, I. Aroso, M. Martins, R. L. Reis and A. R. C. Duarte, *ACS Sustain. Chem. Eng.*, 2014, **2**, 1063–1071.
- 24 S. Khodaverdian, B. Dabirmanesh, A. Heydari, E. Dashtban-Moghadam, K. Khajeh and F. Ghazi, *Int. J. Biol. Macromol.*, 2018, **107**, 2574–2579.
- 25 Y. Ma, Y. Li, S. Ali, P. Li, W. Zhang, M. C. Rauch, S. J. P. Willot, D. Ribitsch, Y. H. Choi and M. Alcalde, *ChemCatChem*, 2020, **12**, 989–994.
- 26 M. Faggian, S. Sut, B. Perissutti, V. Baldan, I. Grabnar and S. Dall'Acqua, *Molecules*, 2016, **21**, 1531.
- 27 H. Vanda, R. Verpoorte, P. G. Klinkhamer and Y. H. Choi, *Deep Eutectic Solvents: Synthesis, Properties, and Applications*, 2019, pp. 61–81.
- 28 A. E. Delorme, J.-M. Andanson and V. Verney, *Int. J. Biol. Macromol.*, 2020, **163**, 919–926.
- 29 A. Kovács, M. Yusupov, I. Cornet, P. Billen and E. C. Neyts, *J. Mol. Liq.*, 2022, **366**, 120180.
- 30 L. Meneses, N. F. Gajardo-Parra, E. Cea-Klapp, J. M. Garrido, C. Held, A. R. Duarte and A. Paiva, *RSC Sustainability*, 2023, **1**, 886–897.
- 31 N. F. Gajardo-Parra, L. Meneses, A. R. C. Duarte, A. Paiva and C. Held, *ACS Sustain. Chem. Eng.*, 2022, **10**, 12873–12881.
- 32 Y. Zhou, Y.-J. Wu, L. Wang, J. Han, J.-C. Wu, C.-M. Li and Y. Wang, *Int. J. Biol. Macromol.*, 2021, **190**, 206–213.
- 33 C. Knudsen, K. Bavishi, K. M. Viborg, D. P. Drew, H. T. Simonsen, M. S. Motawia, B. L. Møller and T. Laursen, *Phytochemistry*, 2020, **170**, 112214.
- 34 M. Mirzaei, H. Khorshahi, H. Tebyanian, R. S. Moradi, M. Rastegar, S. Panahi, R. Sariri and A. M. Latifi, *Biointerface Res. Appl. Chem.*, 2020, **10**, 6488–6497.
- 35 M. S. Lee, K. Lee, M. W. Nam, K. M. Jeong, J. E. Lee, N. W. Kim, Y. Yin, S. Y. Lim, J. Lee and J. H. Jeong, *J. Ind. Eng. Chem.*, 2018, **65**, 343–348.
- 36 Y. Dai, J. van Spronsen, G.-J. Witkamp, R. Verpoorte and Y. H. Choi, *Anal. Chim. Acta*, 2013, **766**, 61–68.
- 37 Y. Tian, M. Zhu, T. Hu and C. Liu, *Int. J. Biol. Macromol.*, 2023, 125477.
- 38 B. S. Chang, K. H. Park and D. B. Lund, *J. Food Sci.*, 1988, **53**, 920–923.
- 39 D. Nash, L. Paleg and J. Wiskich, *Funct. Plant Biol.*, 1982, **9**, 47–57.
- 40 L. Paleg, T. Douglas, A. Van Daal and D. Keech, *Funct. Plant Biol.*, 1981, **8**, 107–114.
- 41 A. Nieter, S. Kelle, M. Takenberg, D. Linke, M. Bunzel, L. Popper and R. G. Berger, *Food Chem.*, 2016, **209**, 1–9.
- 42 D. Linke, A. B. Omarini, M. Takenberg, S. Kelle and R. G. Berger, *Appl. Biochem. Biotechnol.*, 2019, **187**, 894–912.
- 43 P. Bergmann, M. Takenberg, C. Frank, M. Zschätzsch, A. Werner, R. G. Berger and F. Ersoy, *Fermentation*, 2022, **8**, 541.
- 44 P. Bergmann, C. Frank, O. Reinhardt, M. Takenberg, A. Werner, R. G. Berger, F. Ersoy and M. Zschätzsch, *Fermentation*, 2022, **8**, 684.
- 45 K. Shah and R. S. Dubey, *Biol. Plantarum*, 1997, **40**, 121–130.
- 46 X. Liang, L. Zhang, S. K. Natarajan and D. F. Becker, *Antioxidants Redox Signal.*, 2013, **19**, 998–1011.
- 47 B.-P. Wu, Q. Wen, H. Xu and Z. Yang, *J. Mol. Catal. B: Enzym.*, 2014, **101**, 101–107.
- 48 F. Mamashli, J. Badraghi, B. Delavari, H. Lanjanian, M. Sabbaghian, M. Hosseini and A. A. Saboury, *J. Mol. Liq.*, 2018, **272**, 597–608.
- 49 F. Mamashli, J. Badraghi, B. Delavari, M. Sabbaghian, M. Hosseini and A. A. Saboury, *J. Solution Chem.*, 2019, **48**, 689–701.
- 50 T. Singh, D. Aspinall and L. Paleg, *Nat. New Biol.*, 1972, **236**, 188–190.
- 51 R. Storey and R. W. Jones, *Plant Sci. Lett.*, 1975, **4**, 161–168.
- 52 T. M. Chu, D. Aspinall and L. Paleg, *Funct. Plant Biol.*, 1974, **1**, 87–97.
- 53 F. Zulfqar, M. Ashraf and K. H. Siddique, *Agronomy*, 2022, **12**, 276.
- 54 C. De Virgilio, T. Hottiger, J. Dominguez, T. Boller and A. Wiemken, *Eur. J. Biochem.*, 1994, **219**, 179–186.
- 55 A. Wiemken, *Antonie van Leeuwenhoek*, 1990, **58**, 209–217.



- 56 A. A. Elgharbawy, A. Hayyan, M. Hayyan, S. N. Rashid, M. R. M. Nor, M. Y. Zulkifli, Y. Alias and M. E. S. Mirghani, *Chem. Biochem. Eng. Q.*, 2018, **32**, 359–370.
- 57 B. Nian, G. Liao, Y. Song, Y. Su, C. Cao and Y. Liu, *J. Catal.*, 2020, **384**, 159–168.
- 58 H. Monhemi, M. R. Housaindokht, A. A. Moosavi-Movahedi and M. R. Bozorgmehr, *Phys. Chem. Chem. Phys.*, 2014, **16**, 14882–14893.
- 59 F. Thomas and O. Kayser, *Biochem. Eng. J.*, 2022, **180**, 108380.
- 60 L. Yang, Y. Liu, Y. Peng, Y.-C. He and C. Ma, *Catal. Commun.*, 2023, **184**, 106783.
- 61 A. Kovács, E. C. Neyts, I. Cornet, M. Wijnants and P. Billen, *ChemSusChem*, 2020, **13**, 3789–3804.

