

LETTER • OPEN ACCESS

Amino acid production exceeds plant nitrogen demand in Siberian tundra

To cite this article: Birgit Wild *et al* 2018 *Environ. Res. Lett.* **13** 034002

View the [article online](#) for updates and enhancements.

Related content

- [Longer thaw seasons increase nitrogen availability for leaching during fall in tundra soils](#)
Claire C Treat, Wilfred M Wollheim, Ruth K Varner *et al.*
- [Belowground plant biomass allocation in tundra ecosystems and its relationship with temperature](#)
Peng Wang, Monique M P D Heijmans, Liesje Mommer *et al.*
- [Effects of thermo-erosional disturbance on surface soil carbon and nitrogen dynamics in upland arctic tundra](#)
Camila Pizano, Andrés F Barón, Edward A G Schuur *et al.*

Recent citations

- [Divergent patterns of experimental and model-derived permafrost ecosystem carbon dynamics in response to Arctic warming](#)
Christina Schädel *et al*

Environmental Research Letters



LETTER

Amino acid production exceeds plant nitrogen demand in Siberian tundra

OPEN ACCESS

RECEIVED

4 July 2017

REVISED

27 November 2017

ACCEPTED FOR PUBLICATION

4 January 2018

PUBLISHED

16 February 2018

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Birgit Wild^{1,2,3,4,5,17} , Ricardo J Eloy Alves^{2,6}, Jiří Bárta⁷, Petr Čapek⁷, Norman Gentsch⁸, Georg Guggenberger^{8,9}, Gustaf Hugelius^{5,10,11}, Anna Knoltsch^{1,2}, Peter Kuhry¹⁰, Nikolay Lashchinskiy¹², Robert Mikutta¹³, Juri Palmtag¹⁰, Judith Prommer¹, Jörg Schneckner^{1,14}, Olga Shibistova^{8,9}, Mounir Takriti^{1,2,15}, Tim Urich^{2,6,16} and Andreas Richter^{1,2}

¹ Department of Microbiology and Ecosystem Science, University of Vienna, Vienna, Austria

² Austrian Polar Research Institute, Vienna, Austria

³ Department of Earth Sciences, University of Gothenburg, Gothenburg, Sweden

⁴ Department of Environmental Science and Analytical Chemistry, Stockholm University, Stockholm, Sweden

⁵ Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden

⁶ Department of Ecogenomics and Systems Biology, University of Vienna, Vienna, Austria

⁷ Department of Ecosystem Biology, University of South Bohemia, České Budějovice, Czech Republic

⁸ Institute of Soil Science, Leibniz Universität Hannover, Hannover, Germany

⁹ VN Sukachev Institute of Forest, Siberian Branch of Russian Academy of Sciences, Krasnoyarsk, Russia

¹⁰ Department of Physical Geography, Stockholm University, Stockholm, Sweden

¹¹ Department of Earth Science, Stanford University, Stanford, CA, United States of America

¹² Central Siberian Botanical Garden, Siberian Branch of Russian Academy of Sciences, Novosibirsk, Russia

¹³ Soil Science and Soil Protection, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany

¹⁴ Department of Natural Resources and the Environment, University of New Hampshire, Durham, NH, United States of America

¹⁵ Lancaster Environment Centre, Lancaster University, Lancaster, United Kingdom

¹⁶ Institute of Microbiology, Ernst-Moritz-Arndt University, Greifswald, Germany

¹⁷ Author to whom any correspondence should be addressed.

E-mail: birgit.wild@aces.su.se

Keywords: permafrost, tundra, protein depolymerization, nitrogen mineralization, nitrogen limitation, plant productivity

Abstract

Arctic plant productivity is often limited by low soil N availability. This has been attributed to slow breakdown of N-containing polymers in litter and soil organic matter (SOM) into smaller, available units, and to shallow plant rooting constrained by permafrost and high soil moisture. Using ¹⁵N pool dilution assays, we here quantified gross amino acid and ammonium production rates in 97 active layer samples from four sites across the Siberian Arctic. We found that amino acid production in organic layers alone exceeded literature-based estimates of maximum plant N uptake 17-fold and therefore reject the hypothesis that arctic plant N limitation results from slow SOM breakdown. High microbial N use efficiency in organic layers rather suggests strong competition of microorganisms and plants in the dominant rooting zone. Deeper horizons showed lower amino acid production rates per volume, but also lower microbial N use efficiency. Permafrost thaw together with soil drainage might facilitate deeper plant rooting and uptake of previously inaccessible subsoil N, and thereby promote plant productivity in arctic ecosystems. We conclude that changes in microbial decomposer activity, microbial N utilization and plant root density with soil depth interactively control N availability for plants in the Arctic.

Introduction

Arctic plant productivity is limited by low temperatures (Rustad *et al* 2001, Natali *et al* 2012, Sistla *et al* 2013, Leffler *et al* 2016) and low soil N availability (Haag 1974, Shaver and Chapin 1980, Gough *et al* 2002, Gough and Hobbie 2003). Plants meet their N demand

by taking up small compounds such as oligopeptides, amino acids, ammonium, and nitrate (Hill *et al* 2011, Kuzyakov and Xu 2013), whereas the polymers that contain most of the soil N—predominantly proteins, but also heterocyclic compounds (Knicker 2011)—are too large for direct uptake. Nitrogen availability for plants consequently depends on microbial

depolymerization rates (Schimel and Bennett 2004). Nitrogen limitation of arctic plants has been attributed to slow depolymerization under the cold and moist conditions in arctic soils (Hobbie *et al* 2002), and an increase in plant productivity with warming to an increase in soil N availability for plant growth (Chapin 1983, Hartley *et al* 1999, Shaver and Jonasson 1999, Natali *et al* 2012). This perception is supported by a recent modelling study that suggests a stimulation of plant productivity in a warming Arctic by increased N availability in surface soils (Koven *et al* 2015).

Nitrogen uptake by arctic plants is mainly confined to the top 30 cm of the soil where 95% of plant roots are located (Schenk and Jackson 2002, see also Iversen *et al* 2015). Rooting depth is limited by the thickness of the seasonally thawed active layer above the continuously frozen permafrost (Iversen *et al* 2015), and by high soil moisture in the deep active layer (Gebauer *et al* 1996, Iversen *et al* 2015). Permafrost thaw (Vaughan *et al* 2013) and improving drainage (Natali *et al* 2015) with global warming might promote deeper plant rooting and N uptake from previously inaccessible subsoil pools. If subsoils contain relevant amounts of available N, deeper rooting could alleviate plant N limitation, and stimulate plant CO₂ uptake in the Arctic.

Recent inventories show that subsoils contain a large proportion of N in the top meter of arctic permafrost soils, for instance across Siberia, 66%–76% (horizons below 30 cm; Zubrzycki *et al* 2013), 65% (horizons below 30 cm; Palmtag *et al* 2016 and personal communication J. Palmtag), and 81%–89% (Ajj, B, and C horizons; Gentsch *et al* 2015). Subsoils dominate N storage only due to their mass, whereas N concentration per soil mass decreases with depth (Zubrzycki *et al* 2013, Gentsch *et al* 2015), mineral protection of SOM increases (Gentsch *et al* 2015), and consequently protein depolymerization rates decrease (Wild *et al* 2013, 2015). While these observations imply a decrease in the release of potentially plant available N with depth, actual availability also depends on competition with soil microorganisms. Like plants, microorganisms can take up small organic and inorganic N forms, and previous tracer studies show that microorganisms by far outcompete plants for both N forms in arctic organic layers (Schimel and Chapin 1996, Nordin *et al* 2004, Clemmensen *et al* 2008, Lærkedal Sorensen *et al* 2008, Larsen *et al* 2012). With increasing soil depth, however, microbial competition for N might decrease, as indicated by changes in the microbial release of excess N as ammonium into the soil solution ('N mineralization'; Schimel and Bennett 2004). Although gross N mineralization rates typically decrease with soil depth when related to dry soil mass, they have been found to increase when normalized by microbial biomass or microbial N uptake in many systems, including arctic permafrost soils (Mooshammer *et al* 2014, Wild *et al* 2015). This pattern points at an increase in microbial N excess with depth

that contrasts the decrease in protein depolymerization rates, and has been attributed to an even stronger decrease in C availability that constrains microbial growth and N demand (Mooshammer *et al* 2014, Wild *et al* 2015). Subsoil microorganisms might use N-containing organic compounds such as amino acids primarily as C rather than N sources, and release the excess N as ammonium, which thus re-enters the N pool available to plants and microorganisms.

Our study aimed to assess the potential for plant N uptake in different horizons of arctic permafrost soils, and the implications for plant productivity in the case of deeper plant rooting in a future climate. Specifically, we quantified gross production rates of plant available N in different horizons of arctic permafrost soils using ¹⁵N pool dilution assays, and related these rates to estimates of plant N uptake. Active layer horizons were sampled from four upland tundra sites across the Siberian Arctic, comprising 97 samples from 27 soil profiles. We measured gross production rates of amino acids by depolymerization of soil proteins ('protein depolymerization'), and gross release rates of excess N as ammonium from microbial cells ('N mineralization'). In contrast to concentrations of available N pools, gross production rates inform about the replenishment of N pools over time, and thus provide a more useful measure of N availability. As proteins represent the largest soil N pool (Knicker 2011), protein depolymerization is thought to dominate the production of small organic N forms available to plants and microorganisms (Schimel and Bennett 2004, Jan *et al* 2009). We further calculated microbial N use efficiency (NUE) based on gross N transformation rates, as the proportion of N taken up by microorganisms that is used for growth and enzyme synthesis as opposed to mineralization (Mooshammer *et al* 2014). We hypothesized that (1) gross protein depolymerization per soil mass decreases with depth, following a decline in N concentration per soil mass, and that (2) microbial NUE decreases with depth, reflecting a transition from microbial N limitation to microbial N excess. Using soil profile sketches digitized from photographs, we estimated gross amino acid and ammonium production rates in the total active layer per soil surface area, as well as the contribution of individual horizons to the total rates, and compared these values with literature-based estimates of plant N uptake in arctic ecosystems.

Material and methods

Sites and soil sampling

Soils were sampled in the late growing season (August 2010, 2011 or 2012) at four sites across the Siberian Arctic, Cherskiy (69.21°N, 161.69°E), Ari-Mas (72.49°N, 101.66°E), Logata (73.43°N, 98.41°E), and Tazovskiy (67.17°N, 78.91°E; figure 1). Sites are described briefly in table 1, and in detail in Gentsch *et al* (2015). All soils were underlain by continuous permafrost, and




Table 1. Characterization of sampling sites across the Siberian Arctic, including vegetation type, dominant plant species, soil type (Soil Survey Staff 2010), and active layer depth.

	Vegetation type	Dominant plant species	Soil type	Active layer
Cherskiy	Shrubby grass tundra	<i>Betula exilis</i> , <i>Salix spenophylla</i> , <i>Carex lugens</i> , <i>Calamagrostis holmii</i> , <i>Aulacomnium turgidum</i>	Ruptic-Histic Aquiturbel	30–70 cm
	Shrubby tussock tundra	<i>Eriophorum vaginatum</i> , <i>Carex lugens</i> , <i>Betula exilis</i> , <i>Salix pulchra</i> , <i>Aulacomnium turgidum</i>	Ruptic-Histic Aquiturbel	35–60 cm
	Shrubby lichen tundra	<i>Betula exilis</i> , <i>Vaccinium uliginosum</i> , <i>Flavocetraria nivalis</i> , <i>Flavocetraria cucullata</i>	Typic Aquiturbel	35–90 cm
Ari-Mas	Shrubby moss tundra	<i>Betula nana</i> , <i>Dryas punctata</i> , <i>Vaccinium uliginosum</i> , <i>Carex arctisibirica</i> , <i>Aulacomnium turgidum</i>	Typic Aquiturbel	60–85 cm
	Shrubby moss tundra	<i>Cassiope tetragona</i> , <i>Carex arctisibirica</i> , <i>Aulacomnium turgidum</i>	Typic Aquiturbel	65–90 cm
Logata	Dryas tundra	<i>Dryas punctata</i> , <i>Rhytidium rugosum</i> , <i>Hylocomium splendens</i>	Typic Aquiturbel	35–70 cm
	Grassy moss tundra	<i>Betula nana</i> , <i>Carex arctisibirica</i> , <i>Hylocomium splendens</i> , <i>Tomentypnum nitens</i>	Typic Aquiturbel	30–65 cm
Tazovski	Shrubby lichen tundra	<i>Empetrum nigrum</i> , <i>Ledum palustre</i> , <i>Betula nana</i> , <i>Cladonia rangiferina</i> , <i>Cladonia stellaris</i>	Typic Aquiturbel	100–120 cm
	Forest tundra	<i>Larix sibirica</i> , <i>Ledum palustra</i> , <i>Betula nana</i> , <i>Vaccinium uliginosum</i> , <i>Cladonia rangiferina</i> , <i>Cladonia stellaris</i>	Typic Aquiturbel	130–150 cm

described as Typic or Ruptic-Histic Aquiturbels (Soil Survey Staff 2010).

We selected two or three representative upland tundra vegetation types per site and in each excavated three 5 m long soil trenches down to the permafrost table. We described all active layer horizons at high resolution, sampled them individually, and determined their bulk density (for details see Gentsch *et al* 2015). Living roots were removed immediately and samples were stored at surface soil temperature for a maximum of a few days before analysis. Samples were not sieved, but carefully homogenized by hand.

We categorized samples into four horizon classes as in Wild *et al* (2016), namely organic layers (O horizons), mineral topsoils (OA, A, and AB horizons), mineral subsoils (B, BC, and C horizons), and subducted topsoils, i.e. pockets of poorly decomposed

topsoil material subducted into the subsoil by freeze-thaw processes (Ojj, OAjj, and Ajj horizons; Ping *et al* 2015). Based on soil profile sketches digitized from photographs, we calculated soil volumes per soil surface area for each pedon and horizon class as described by Gentsch *et al* (2015). A subset of soil samples that covered all depths and all four sites ($n = 97$) was used to measure soil C and N pools and gross N transformation rates. pH values were measured in aqueous suspension at a solid:solution ratio of 1:2.5.

Carbon and nitrogen pools

Total soil organic C and N, as well as ^{13}C and ^{15}N content were measured in dried and ground samples using elemental analysis-isotope ratio mass spectrometry (EA-IRMS; Elementar vario MICRO cube EA, Elementar Analysensysteme GmbH, Germany, and

Table 2. Characterization of the sampled soil horizons. Values represent means \pm standard errors; significant differences between soil horizon categories are indicated by different letters ($p < 0.05$). 'Depth' refers to mean sample depth for each horizon class.

	Number	Depth (cm)	Organic C (%)	N (%)	C/N (w/w)	$\delta^{13}\text{C}$ (‰)	$\delta^{15}\text{N}$ (‰)	pH
Org. layer	19	9 \pm 1a	22.98 \pm 1.67a	0.90 \pm 0.05a	25.83 \pm 1.34a	-28.11 \pm 0.19c	0.82 \pm 0.31c	5.41 \pm 0.15c
Min. topsoil	22	16 \pm 2b	4.18 \pm 0.47b	0.26 \pm 0.03c	16.35 \pm 0.62b	-27.29 \pm 0.18b	2.78 \pm 0.25b	5.95 \pm 0.15b
Subd. topsoil	38	43 \pm 3c	6.77 \pm 0.66b	0.40 \pm 0.03b	16.16 \pm 0.47b	-27.18 \pm 0.12b	2.69 \pm 0.29b	6.26 \pm 0.11ab
Min. subsoil	18	49 \pm 8c	1.14 \pm 0.18c	0.09 \pm 0.01d	11.62 \pm 0.57c	-25.70 \pm 0.30a	4.99 \pm 0.35a	6.50 \pm 0.14a

Table 3. Microbial C and N, dissolved organic C (DOC), dissolved organic N (DON), ammonium and nitrate concentrations in the sampled soil horizons. Values represent means \pm standard errors related to dry soil (d.s.); significant differences between soil horizon categories are indicated by different letters ($p < 0.05$). Microbial C and N were not analyzed in the Cherskiy samples and are not included here.

	Microbial C ($\mu\text{g C g}^{-1}$ d. s.)	Microbial N ($\mu\text{g N g}^{-1}$ d. s.)	DOC ($\mu\text{g C g}^{-1}$ d. s.)	DON ($\mu\text{g N g}^{-1}$ d. s.)	Ammonium ($\mu\text{g N g}^{-1}$ d. s.)	Nitrate ($\mu\text{g N g}^{-1}$ d. s.)
Org. layer	936.8 \pm 276.6a	103.9 \pm 19.8a	546.2 \pm 259.8a	52.77 \pm 35.04a	5.67 \pm 1.25a	2.17 \pm 0.68ab
Min. topsoil	401.6 \pm 96.1a	36.5 \pm 6.6b	86.1 \pm 12.9b	5.99 \pm 1.74b	1.30 \pm 0.38b	1.25 \pm 0.52b
Subd. topsoil	114.1 \pm 18.7b	10.7 \pm 2.1c	68.6 \pm 7.5b	3.29 \pm 0.76c	1.97 \pm 0.33b	2.25 \pm 0.42a
Min. subsoil	47.5 \pm 11.7c	2.9 \pm 0.7d	26.6 \pm 4.8c	0.75 \pm 0.32d	2.60 \pm 0.85b	0.80 \pm 0.28b

Elementar IsoPrime 100 IRMS, IsoPrime Ltd, UK), and C/N ratios were calculated on a mass basis. Data are presented in table 2. Concentrations of dissolved organic C (DOC) and total dissolved N were measured in 0.5 M K_2SO_4 extracts (Cherskiy, Ari-Mas, Logata) or 1 M KCl extracts (Tazovskiy) using a TOC/TN analyzer (Cherskiy, Ari-Mas, Logata: Elementar LiquiC/TOC II; Tazovskiy: Shimadzu TOC-VCPH/CPN/TNM-1). Concentrations of ammonium and nitrate were measured photometrically in 1 M KCl extracts following Kandeler and Gerber (1988) and Miranda *et al* (2001), respectively, and concentrations of dissolved organic N (DON) were calculated by subtracting ammonium and nitrate from total dissolved N. Microbial C and N were estimated using the chloroform fumigation method (Brookes *et al* 1985). Aliquots of fresh soil were fumigated with ethanol-free chloroform, dissolved organic C and total dissolved N were extracted and analyzed as described above, and microbial C and N were calculated as the difference between fumigated and non-fumigated samples. We did not apply a correction factor to account for incomplete recovery of microbial C and N (Brookes *et al* 1985). Microbial C and N were not determined in the Cherskiy samples. Data on microbial C and N, DOC, DON, ammonium and nitrate are presented in table 3.

Gross nitrogen transformation rates

Gross rates of protein depolymerization, microbial amino acid uptake, N mineralization, and microbial ammonium uptake were measured using ^{15}N pool dilution assays (Kirkham and Bartholomew 1954). Briefly, we labelled the amino acid (protein depolymerization and amino acid uptake) or ammonium pool (N mineralization and ammonium uptake) with ^{15}N , determined concentration and isotopic composition of the respective pool at two time points (amino acids: 10 and 30 min; ammonium: 4 and 24 h), and calculated the fluxes into and out of the pool from the dilution of ^{15}N and changes in concentration between the two time points.

Gross protein depolymerization and amino acid uptake rates were determined following Wanek *et al* (2010), as modified by Wild *et al* (2013). Duplicates of fresh soil (1–4 g depending on SOM content) were amended with ^{15}N labelled amino acids (mixture of 20 amino acids with $>98\%$ ^{15}N , Spectra and Cambridge Isotope Laboratories, 0.31–2.5 μg per g fresh soil adjusted to the expected amino acid content of the sample). Sample duplicates were extracted after 10 or 30 min, respectively, with 20 ml 10 mM CaSO_4 containing 3.7% formaldehyde. Samples were filtered, loaded on pre-cleaned cation exchange cartridges (OnGuard II H 1 cc cartridges, Thermo/Dionex), eluted with 3 M NH_3 , amended with internal standards (nor-valine, nor-leucine, and para-chloro-phenylalanine, 1 μg each per sample, Sigma-Aldrich), dried, derivatized with ethyl-chloroformate (Wanek *et al* 2010), and analyzed with gas chromatography-mass spectrometry (Thermo Trace GC Ultra with Thermo DSQ II or ISQ mass spectrometer). Blanks and one set of amino acid standards were processed with the samples throughout the procedure to correct for incomplete recovery. A second set of amino acid standards was derivatized with the samples for calibration of alanine, isoleucine, leucine, phenylalanine, proline, valine, aspartate/asparagine, and glutamate/glutamine concentrations. Amino acid ^{15}N contents were calculated from peak areas of light and heavy fragments as described by Wanek *et al* (2010).

Gross N mineralization and ammonium uptake rates were determined as in Wild *et al* (2013), by amending duplicates of fresh soil (1–4 g depending on SOM content) with 500 μl of 0.125 mM ^{15}N labelled $(\text{NH}_4)_2\text{SO}_4$ (10 at%, Sigma-Aldrich). Sample duplicates were extracted after 4 or 24 h, respectively, with 13 ml 2 M KCl, and filtered through ash free filter paper. Ammonium was converted into ammonia which diffused into acid traps (Sørensen and Jensen 1991), and amount and isotopic composition were measured with EA-IRMS (CE Instrument EA 1110 elemental analyzer, Finnigan MAT ConFlo II interface, Finnigan MAT DeltaPlus IRMS).

The calculation of gross rates based on pool dilution assays requires an enrichment of ^{15}N in the target pool (here: amino acids and ammonium, respectively) at both time points, and a depletion of ^{15}N between them. This was confirmed for all samples. The amino acid pool showed average ^{15}N contents of $26.4 \pm 2.5 \text{ at\%}$ (mean \pm standard error) at time point 1 and $15.0 \pm 1.8 \text{ at\%}$ at time point 2 after addition of amino acids with $> 98 \text{ at\% } ^{15}\text{N}$, and the ammonium pool of $2.3 \pm 0.2 \text{ at\%}$ at time point 1 and $1.2 \pm 0.1 \text{ at\%}$ at time point 2 after addition of $(\text{NH}_4)_2\text{SO}_4$ with $10 \text{ at\% } ^{15}\text{N}$. Gross rates of protein depolymerization, microbial amino acid uptake, N mineralization, and microbial ammonium uptake were calculated following Kirkham and Bartholomew (1954), and normalized by dry soil mass, soil N, and soil volume using bulk density values of the respective samples. We then estimated gross N transformation rates per soil surface area, by averaging gross rates per soil volume for each horizon class, and multiplying with the average soil volume per soil surface area of the respective horizon class. Standard errors were calculated following Gaussian error propagation.

We further estimated microbial N use efficiency (NUE) as the proportion of N taken up that was not mineralized, but used for microbial growth and enzyme synthesis. Previous studies on microbial NUE have considered only microbial amino acid uptake as it dominated microbial N uptake in their samples (Wild *et al* 2013, Mooshammer *et al* 2014). We extended their equation to include ammonium uptake that especially in mineral subsoils contributed considerably to microbial N uptake. Microbial NUE was calculated from gross rates of microbial amino acid and ammonium uptake as well as N mineralization as:

$$\text{NUE} = \frac{\text{Amino acid and ammonium uptake} - \text{ammonium release}}{\text{Amino acid and ammonium uptake}}. \quad (1)$$

Given the limited infrastructure at our remote field sites in the Siberian Arctic, we measured all gross N transformation rates at surface soil temperatures, and thereby possibly overestimated rates in the deeper soil. We did not apply a temperature correction to the measured rates, but estimated the magnitude of the temperature effect on gross N transformation rates in the deeper soil. Cherskiy data are partly presented in Wild *et al* (2013).

Estimate of plant nitrogen uptake

We estimated maximum plant N uptake in arctic ecosystems based on net primary production rates reported in previous studies. For the Canadian Arctic, total net primary production rates from less than $20 \text{ g m}^{-2} \text{ yr}^{-1}$ in polar deserts to up to $1000 \text{ g m}^{-2} \text{ yr}^{-1}$ in highly productive shrub ecosystems have been suggested (Gould *et al* 2003), and this range likely applies also to other areas across the Arctic. Total net


primary production rates of $200\text{--}300 \text{ g m}^{-2} \text{ yr}^{-1}$ have been observed in heath tundra in the Scandinavian sub-Arctic (Campioli *et al* 2009, Jonasson *et al* 1999), and aboveground net primary production rates of $30\text{--}400 \text{ g m}^{-2} \text{ yr}^{-1}$ at sites dominated by heath shrubs, tall deciduous shrubs or tussock-forming graminoids in the Alaskan Arctic (Shaver and Chapin 1991, Schuur *et al* 2007, Natali *et al* 2012, DeMarco *et al* 2014). Considering that belowground production likely represents only a small fraction of total production (e.g. 15% in a sub-Arctic heath tundra; Campioli *et al* 2009), these rates fall well into the range of total net primary production presented above. For our estimate of maximum plant N uptake, we therefore assumed maximum net primary production of $1000 \text{ g m}^{-2} \text{ yr}^{-1}$, 1% N content in the new biomass (Jonasson *et al* 1999, Campioli *et al* 2009), 120 growing season days (Ernakovich *et al* 2014), and constant plant N uptake during the growing season (see discussion). We thus arrived at a maximum daily plant N uptake of $83 \text{ mg N m}^{-2} \text{ d}^{-1}$ during the growing season (total range $1.7\text{--}83 \text{ mg N m}^{-2} \text{ d}^{-1}$). Our estimate of maximum plant N uptake by far exceeds previous estimates of actual plant N uptake of $1 \text{ g N m}^{-2} \text{ yr}^{-1}$ ($8.3 \text{ mg N m}^{-2} \text{ d}^{-1}$ assuming 120 growing season days) in a sub-arctic heath tundra (Jonasson *et al* 1999), and of 0.8 and $1.5 \text{ g N m}^{-2} \text{ yr}^{-1}$ (6.3 and $12.2 \text{ mg N m}^{-2} \text{ d}^{-1}$) in tussock tundra (only aboveground biomass considered; Chapin *et al* 1988). We are therefore confident that our maximum value represents a conservative upper limit for plant N uptake, as it might be achieved in the most productive arctic ecosystems.

Statistical analyses

We tested for significant differences ($p < 0.05$) between horizon classes using ANOVA with Tukey HSD as post hoc test, after rank-transformation where necessary to meet the conditions. Where conditions could not be met even after transformation, we conducted non-parametric Mann-Whitney-U tests. All statistics were performed in R 3.1.3 (R Development Core Team 2015), with the additional package GenABEL (Aulchenko *et al* 2007).

Results

Gross protein depolymerization rates significantly decreased with depth, from organic layers to mineral topsoils and further to subducted topsoils and mineral subsoils (figure 2). This pattern was observed when rates were related to dry soil mass, but also when rates were normalized by soil N content, although less pronounced. In support of hypothesis (1), our data thus suggest that the decrease in protein depolymerization with depth was driven by a combination of decreasing N concentration per soil mass (i.e. substrate availability; table 2), and decreasing microbial depolymerization activity per soil N. Although mineral




and subducted topsoils have similar SOM properties (table 2; see also Xu *et al* 2009), subducted topsoil rates accounted for only 37% of mineral topsoil rates when related to soil N. A deceleration of N transformations by subduction has been previously described (Wild *et al* 2013) and likely reflects a general deceleration of microbial decomposition processes (Kaiser *et al* 2007, Čapek *et al* 2015, Wild *et al* 2016).

Gross N mineralization rates decreased with depth only when related to dry soil mass, but were in the same range in all horizon classes when normalized by N content. In support of hypothesis (2), the different patterns of protein depolymerization and

N mineralization rates were reflected in a significant decrease in microbial NUE with depth, from 0.92 ± 0.02 (mean \pm standard error) in organic layers to 0.81 ± 0.03 in mineral topsoils, 0.67 ± 0.04 in subducted topsoils, and 0.61 ± 0.07 in mineral subsoils (figure 3). Microorganisms in organic layers used 92% of the N taken up for growth and enzyme synthesis, and mineralized 8%, whereas microorganisms in mineral subsoils used 61%, and mineralized 39%.

Using bulk density values of the sampled soils, we calculated N transformation rates per soil volume, and thereby normalized the amount of potentially plant available N by the volume plant roots need to occupy for its uptake. Also related to soil volume, gross



protein depolymerization rates significantly decreased with depth (figure 2), and gross N mineralization rates were in the same range in all horizons. We then used digitized soil profile sketches to calculate the thickness of different active layer horizons, and their volume within the active layer per soil surface area. Given that mineral subsoils represented 78% of the active layer volume at our study sites, they dominated both protein depolymerization and N mineralization rates per soil surface area, contributing 46% and 68%, respectively, to the total active layer rates (figure 4). Organic layers contributed 32% to protein depolymerization, but only 7% to N mineralization, mineral topsoils 12% and 10%, and subducted topsoils 10% and 14%, respectively.

We note that gross rates in deeper soils might be overestimated as they were measured at surface soil temperatures. During our sampling campaigns, soil temperatures decreased from 8.3 °C at the average organic layer depth to 7.1 °C in mineral topsoils, 4.4 °C in subducted topsoils, and 3.8 °C in mineral subsoils. Assuming a Q_{10} of 2, subducted topsoil rates would be lower by 24% under *in situ* temperatures, and mineral subsoil rates by 27%. Organic layers would then contribute 38% and 10%, mineral topsoils 13% and 12%, subducted topsoils 9% and 14%, and mineral subsoils 40% and 65% to protein depolymerization and N mineralization, respectively, in the active layer.

Discussion


Arctic plant productivity is often limited by low soil N availability (Haag 1974, Shaver and Chapin 1980, Gough *et al* 2002, Hobbie *et al* 2002, Gough and Hobbie 2003). Based on previous reports of net primary production, we estimate that plant N uptake can reach up to 83 mg N m⁻² d⁻¹ in the most productive arctic ecosystems during the growing season. Even this

maximum value of plant uptake was exceeded by more than an order of magnitude by the release of amino acid N from soil proteins (figure 4). Gross protein depolymerization rates at our study sites amounted to 4.3 ± 0.8 g N m⁻² d⁻¹ (mean \pm standard error) in the entire active layer, and 1.4 ± 0.3 g N m⁻² d⁻¹ in organic layers where most plant roots are located (Schenk and Jackson 2002). Since arctic plants can directly use amino acids as N sources (Schimel and Chapin 1996, Nordin *et al* 2004, Lærkedal Sorensen *et al* 2008), our findings contradict the hypothesis that arctic plant N limitation results from slow SOM breakdown.

We estimated maximum daily plant N uptake under the assumption of constant uptake over the growing season because studies at high time resolution are missing, and we measured gross protein depolymerization rates at one time point in the late growing season. Both parameters likely show seasonal variation, as observed also for microbial biomass and extracellular enzyme activities (Weintraub and Schimel 2005, Wallenstein *et al* 2009, Edwards and Jefferies 2013, Sistla and Schimel 2013, Stark and Väisänen 2014). We expect maximum protein depolymerization rates in the late growing season when soils are warmest, whereas peaks in plant N uptake have been suggested in the late, but also in the early growing season (Olsrud and Christensen 2004, Blume-Werry *et al* 2016, Sloan *et al* 2016). A temporal mismatch between N release from SOM polymers and plant N uptake might thus contribute to constraining arctic plant productivity. Furthermore, gross N transformation rates were measured in homogenized soil samples, which likely show some deviation from rates in undisturbed soils. However, given that average amino acid production rates exceeded even the maximum plant N uptake by more than an order of magnitude, and that amino acids represent only part of the available N pool (see e.g. for oligopeptides Hill *et al* 2011), the discrepancy between soil N availability and plant N uptake seems too large to be attributed to seasonal fluctuations and methodological artefacts alone.

While we did not measure rooting depth at our study sites, previous studies agree that most tundra roots are located in the shallow soil. A meta-analysis reported 50% of tundra roots in the top 9 cm, and 95% in the top 29 cm (Schenk and Jackson 2002), and more recent studies 50% (Alaskan tussock tundra; van Wijk *et al* 2003) and 68%–100% in the top 10 cm (Alaskan tussock-shrub tundra; Zhu *et al* 2016), 46%–65% in the top 5 cm (Siberian tussock-shrub tundra; Wang *et al* 2016), and 78% in the top 2 cm (Svalbard semi-desert; Oulehle *et al* 2016, see also Iversen *et al* 2015 for a recent review). These patterns are in line with our personal observations at our study sites.

Comparing microbial N dynamics in different horizons with typical plant rooting depths suggests that arctic plant N limitation is driven by strong competition with microbes for available N in the dominant rooting zone. Protein depolymerization rates in organic layers



exceeded maximum plant N uptake by a factor of 17 (figure 4), but our data also show that a large proportion of the N made available by protein depolymerization in organic layers was rapidly sequestered within the microbial biomass, as microorganisms retained 92% of the N taken up, and released only 8% by mineralization (figure 3). High microbial NUE indicates high microbial N demand compared to N availability (Mooshammer *et al* 2014), and suggests that under natural conditions, microorganisms compete with plants for available N in organic layers. These conclusions are supported by previous studies in arctic and subarctic ecosystems that reported a by far more efficient incorporation of ¹⁵N added to organic layers into microbial than plant biomass (Schimel and Chapin 1996, Nordin *et al* 2004, Clemmensen *et al* 2008, Lærkedal Sorensen *et al* 2008, Larsen *et al* 2012). A recent global meta-analysis suggests that this pattern might be strongest for organic N (Kuzyakov and Xu 2013) which dominates the available N pool in high latitude organic layers (Jones and Kielland 2002, Larsen *et al* 2012, Wild *et al* 2013, 2015).

Mineral subsoils and subducted topsoils together contributed 56% to gross protein depolymerization and 83% to gross N mineralization in the active layer at our sites. In both horizons, microorganisms exhibited lower NUE than in organic layers (figure 3), indicating higher microbial N excess. With plant N uptake from subsoils currently impeded by shallow rooting, available subsoil N might fuel nitrification/denitrification processes that can promote ecosystem N losses by nitrate leaching and outgassing of N₂ or N₂O (Cameron *et al* 2013). Although gross nitrification rates per microbial biomass have been found to increase with depth in arctic soils (Wild *et al* 2015), and nitrate has been

identified as the dominant N form in artificial leachates of arctic soil samples (Treat *et al* 2016), nitrate leaching under natural conditions is likely constrained by the presence of permafrost. In the discontinuous permafrost zone, however, high nitrate concentrations have been observed in rivers (Petronne *et al* 2006, Cai *et al* 2008), and the subsoil has been suggested as its source (Petronne *et al* 2006). An accumulation of inorganic N, as well as high net N mineralization rates upon thaw have also been observed in the current upper permafrost (Keuper *et al* 2012, Beermann *et al* 2017).

Subsoil N cycling might be substantially altered by changes in temperature, hydrological regime and vegetation composition in the next decades. Active layer deepening (Vaughan *et al* 2013) and improving drainage (Natali *et al* 2015) might permit deeper plant rooting and uptake of subsoil N. Arctic warming could thereby stimulate plant productivity even without accelerating depolymerization or N mineralization rates. However, warmer and drier conditions will likely promote microbial growth and activity in subsoils, consequently stimulate both microbial production and consumption of available N, and enhance competition for N. Increased root density, and thus increased input of plant-derived organic compounds, could additionally stimulate microbial activity and alter microbial N cycling, but also accelerate the mineralization of soil organic C to CO₂ ('priming effect'; Bengtson *et al* 2012, Dijkstra *et al* 2013, Zhu *et al* 2014). Mineral subsoils of arctic permafrost soils are particularly susceptible to plant-derived compounds in laboratory experiments (Wild *et al* 2014, 2016), and we suggest that deeper plant rooting could accelerate soil organic C losses from the deep active layer, and counteract the increased plant CO₂ uptake, as indicated also in

a long-term tundra warming study (Sistla *et al* 2013). This effect, however, might be restricted to sites dominated by deep-rooting plants such as graminoids that can take up N from the subsoil (Oulehle *et al* 2016, Zhu *et al* 2016). Although graminoid abundance might increase at some, especially colder sites within the tundra biome, most sites are expected to experience a decrease in graminoid, and increase in shrub abundance (Elmendorf *et al* 2012). Shrubs take up N from shallower depths than graminoids (Oulehle *et al* 2016, Zhu *et al* 2016) which could effectively reduce plant N uptake from the subsoil, and promote N losses. Systems in the south of the tundra biome might be particularly affected by nitrate leaching as both increases in shrub abundance (Elmendorf *et al* 2012) and decreases in permafrost extent (Chadburn *et al* 2017) have been predicted for these sites.

Conclusions

Comparing N transformations across active layer horizons in the Siberian Arctic, we demonstrate changes in microbial N dynamics with soil depth that, through interactions with plant rooting patterns, might play a crucial role for ecosystem productivity and C storage. We found that production rates of available N greatly exceeded literature-based estimates of plant N uptake. These findings do not contradict previous evidence for N limitation of arctic plant productivity, but show that arctic plant N limitation does not result from slow breakdown of SOM polymers. Instead, we suggest that plant N uptake was constrained by strong microbial N demand in the shallow soil where most plant roots are located. Rising temperatures and improving soil drainage might facilitate plant N uptake from previously inaccessible subsoil horizons, thereby stimulate plant productivity, and reduce N losses in systems with deep-rooting plants. An increasing abundance of shallow-rooting shrubs, as predicted for large areas across the Arctic, could counteract this effect and promote N losses. We conclude that interactions between changes in soil abiotic conditions and vegetation composition could substantially transform C and N cycles in arctic ecosystems, with potential consequences for the global C balance.

Acknowledgments

This study is part of the CryoCARB project (Long-term Carbon Storage in Cryoturbated Arctic Soils), co-funded by the Austrian Science Fund (FWF): I370-B17, the German Federal Ministry of Education and Research (03F0616A), the Czech Ministry of Education, Youth and Sports (MSM 7E10073—CryoCARB), the Russian Ministry of Education and Science (No. 14.B25.31.0031), the Swedish Research Council (824–2009-77357), and the Norwegian Research Fund (NFR): NFR-200411, and was

further supported by a JPI Climate Project (COUP-Austria; BMWFW-6.020/0008) awarded to Andreas Richter. Jiří Bárta and Tim Urich received additional funding from the Czech Science Foundation (16–18453S).

ORCID iDs

Birgit Wild  <https://orcid.org/0000-0002-9611-0815>

References

- Aulchenko Y S, Ripke S, Isaacs A and van Duijn C M 2007 GenABEL: an R library for genome-wide association analysis *Bioinformatics* **23** 1294–6
- Beeremann F, Langer M, Wetterich S, Strauss J, Boike J, Fiencke C, Schirrmeister L, Pfeiffer E-M and Kutzbach L 2017 Permafrost thaw and liberation of inorganic nitrogen in eastern Siberia *Permafrost Periglac.* **28** 605–618
- Bengtson P, Barker J and Grayston S J 2012 Evidence of a strong coupling between root exudation, C and N availability, and stimulated SOM decomposition caused by rhizosphere priming effects *Ecol. Evol.* **2** 1843–52
- Bivand R, Keitt T and Rowlingson B 2015 rgdal: Bindings for the geospatial data abstraction library R package version 0.9–2
- Blume-Werry G, Wilson S D, Kreyling J and Milbau A 2016 The hidden season: growing season is 50% longer below than above ground along an arctic elevation gradient *New Phytol.* **209** 978–86
- Brookes P C, Landman A, Pruden G and Jenkinson D S 1985 Chloroform fumigation and the release of soil nitrogen: a rapid direct extraction method to measure microbial biomass nitrogen in soil *Soil Biol. Biochem.* **17** 837–42
- Cai Y, Guo L, Douglas T A and Whitley T E 2008 Seasonal variations in nutrient concentrations and speciation in the Chena River, Alaska *J. Geophys. Res. Biogeosci.* **113** 1–11
- Cameron K C, Di H J and Moir J L 2013 Nitrogen losses from the soil/plant system: a review *Ann. Appl. Biol.* **162** 145–73
- Campioli M, Michelsen A, Demey A, Vermeulen A, Samson R and Lemeur R 2009 Net primary production and carbon stocks for subarctic mesic-dry tundras with contrasting microtopography, altitude, and dominant species *Ecosystems* **12** 760–76
- Čapek P *et al* 2015 The effect of warming on the vulnerability of subducted organic carbon in arctic soils *Soil Biol. Biochem.* **90** 19–29
- Chadburn S E, Burke E J, Cox P M, Friedlingstein P, Hugelius G and Westermann S 2017 An observation-based constraint on permafrost loss as a function of global warming *Nat. Clim. Change* **7** 340–4
- Chapin F S 1983 Direct and indirect effects of temperature on arctic plants *Polar Biol.* **2** 47–52
- Chapin F S, Fetcher N, Kielland K, Everett K R and Linkings A E 1988 Productivity and nutrient cycling of Alaskan Tundra: enhancement by flowing soil water *Ecology* **69** 693–702
- Clemmensen K E, Sorensen P L, Michelsen A, Jonasson S and Ström L 2008 Site-dependent N uptake from N-form mixtures by arctic plants, soil microbes and ectomycorrhizal fungi *Oecologia* **155** 771–83
- DeMarco J, Mack M C, Bret-Harte M S, Burton M and Shaver G R 2014 Long-term experimental warming and nutrient additions increase productivity in tall deciduous shrub tundra *Ecosphere* **5** 1–22
- Dijkstra F, Carrillo Y, Pendall E and Morgan J A 2013 Rhizosphere priming: a nutrient perspective *Front. Microbiol.* **4** 216
- Edwards K A and Jefferies R L 2013 Inter-annual and seasonal dynamics of soil microbial biomass and nutrients in wet and dry low-Arctic sedge meadows *Soil Biol. Biochem.* **57** 83–9

- Elmendorf S C *et al* 2012 Global assessment of experimental climate warming on tundra vegetation: heterogeneity over space and time *Ecol. Lett.* **15** 164–75
- Ernakovich J G, Hopping K A, Berdanier A B, Simpson R T, Kachergis E J, Steltzer H and Wallenstein M D 2014 Predicted responses of arctic and alpine ecosystems to altered seasonality under climate change *Glob. Change Biol.* **20** 3256–69
- Gebauer R L E, Tenhunen J D and Reynolds J F 1996 Soil aeration in relation to soil physical properties, nitrogen availability, and root characteristics within an arctic watershed *Plant Soil* **178** 37–48
- Gentsch N *et al* 2015 Storage and transformation of organic matter fractions in cryoturbated permafrost soils across the Siberian Arctic *Biogeosciences* **12** 4525–42
- Gough L and Hobbie S E 2003 Responses of moist non-acidic arctic tundra to altered environment: productivity, biomass, and species richness *Oikos* **103** 204–16
- Gough L, Wookey P A and Shaver G R 2002 Dry heath arctic tundra responses to long-term nutrient and light manipulation *Arctic, Antarct. Alp. Res.* **34** 211–8
- Gould W A, Reynolds M and Walker D A 2003 Vegetation, plant biomass, and net primary productivity patterns in the Canadian Arctic *J. Geophys. Res.* **108** 8167
- Haag R W 1974 Nutrient limitations to plant production in two tundra communities *Can. J. Bot.* **52** 103–16
- Hartley A E, Neill C, Melillo J M, Crabtree R and Bowles F P 1999 Plant performance and soil nitrogen mineralization in response to simulated climate change in subarctic dwarf shrub heath *Oikos* **86** 331–43
- Hill P W, Farrar J, Roberts P, Farrell M, Grant H, Newsham K K, Hopkins D W, Bardgett R D and Jones D L 2011 Vascular plant success in a warming Antarctic may be due to efficient nitrogen acquisition *Nat. Clim. Change* **1** 50–3
- Hobbie S E, Nadelhoffer K J and Högberg P 2002 A synthesis: the role of nutrients as constraints on carbon balances in boreal and arctic regions *Plant Soil* **242** 163–70
- Iversen C M, Sloan V L, Sullivan P F, Euskirchen E S, McGuire A D, Norby R J, Walker A P, Warren J M and Wullschlegel S D 2015 The unseen iceberg: plant roots in arctic tundra *New Phytol.* **205** 34–58
- Jan M T, Roberts P, Tonheim S K and Jones D L 2009 Protein breakdown represents a major bottleneck in nitrogen cycling in grassland soils *Soil Biol. Biochem.* **41** 2272–82
- Jonasson S, Michelsen A and Schmidt I K 1999 Coupling of nutrient cycling and carbon dynamics in the Arctic, integration of soil microbial and plant processes *Appl. Soil Ecol.* **11** 135–46
- Jones D L and Kielland K 2002 Soil amino acid turnover dominates the nitrogen flux in permafrost-dominated taiga forest soils *Soil Biol. Biochem.* **34** 209–19
- Kaiser C, Meyer H, Biasi C, Rusalimova O, Barsukov P and Richter A 2007 Conservation of soil organic matter through cryoturbation in arctic soils in Siberia *J. Geophys. Res. Biogeosci.* **112** 1–8
- Kandeler E and Gerber H 1988 Short-term assay of soil urease activity using colorimetric determination of ammonium *Biol. Fertil. Soils* **6** 68–72
- Keuper F, van Bodegom P M, Dorrepaal E, Weedon J T, van Hal J, van Logtestijn R S P and Aerts R 2012 A frozen feast: thawing permafrost increases plant-available nitrogen in subarctic peatlands *Glob. Change Biol.* **18** 1998–2007
- Kirkham D and Bartholomew W V 1954 Equations for following nutrient transformations in soil, utilizing tracer data *Soil Sci. Soc. Am. J.* **18** 33–4
- Knicker H 2011 Soil organic N—an under-rated player for C sequestration in soils? *Soil Biol. Biochem.* **43** 1118–29
- Koven C D, Lawrence D M and Riley W J 2015 Permafrost carbon-climate feedback is sensitive to deep soil carbon decomposability but not deep soil nitrogen dynamics *Proc. Natl Acad. Sci. USA* **112** 3752–7
- Kuzyakov Y and Xu X 2013 Competition between roots and microorganisms for nitrogen: mechanisms and ecological relevance *New Phytol.* **198** 656–69
- Lærkedal Sørensen P, Engelbrecht Clemmensen K, Michelsen A, Jonasson S and Ström L 2008 Plant and microbial uptake and allocation of organic and inorganic nitrogen related to plant growth forms and soil conditions at two subarctic tundra sites in Sweden *Arct. Antarct. Alp. Res.* **1** 171–80
- Larsen K S, Michelsen A, Jonasson S, Beier C and Grogan P 2012 Nitrogen uptake during fall, winter and spring differs among plant functional groups in a subarctic heath ecosystem *Ecosystems* **15** 927–39
- Leffler A J, Klein E S, Oberbauer S F and Welker J M 2016 Coupled long-term summer warming and deeper snow alters species composition and stimulates gross primary productivity in tussock tundra *Oecologia* **181** 287–97
- Lemon J 2006 plotrix: a package in the red light district of R *R-News* **6** 8–12
- Miranda K M, Espey M G and Wind D A 2001 A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite *Nitric Oxide-Biol. Ch.* **5** 62–71
- Mooshammer M *et al* 2014 Adjustment of microbial nitrogen use efficiency to carbon: nitrogen imbalances regulates soil nitrogen cycling *Nat. Commun.* **5** 3694
- Natali S M *et al* 2015 Permafrost thaw and soil moisture driving CO₂ and CH₄ release from upland tundra *J. Geophys. Res. Biogeosci.* **120** 525–37
- Natali S M, Schuur E A G and Rubin R L 2012 Increased plant productivity in Alaskan tundra as a result of experimental warming of soil and permafrost *J. Ecol.* **100** 488–98
- Nordin A, Schmidt I K and Shaver G R 2004 Nitrogen uptake by arctic soil microbes and plants in relation to soil nitrogen supply *Ecology* **85** 955–62
- Olsrud M and Christensen T R 2004 Carbon cycling in subarctic tundra: seasonal variation in ecosystem partitioning based on *in situ* ¹⁴C pulse-labelling *Soil Biol. Biochem.* **36** 245–53
- Oulehle F, Rowe E C, Myška O, Chuman T and Evans C D 2016 Plant functional type affects nitrogen use efficiency in high-Arctic tundra *Soil Biol. Biochem.* **94** 19–28
- Palmtag J, Ramage J, Hugelius G, Gentsch N, Lashchinskiy N, Richter A and Kuhry P 2016 Controls on the storage of organic carbon in permafrost soil in northern Siberia *Eur. J. Soil Sci.* **67** 478–91
- Pebesma E J and Bivand R S 2005 Classes and methods for spatial data in R *R-News* **5** 9–15
- Petrone K C, Jones J B, Hinzman L D and Boone R D 2006 Seasonal export of carbon, nitrogen, and major solutes from Alaskan catchments with discontinuous permafrost *J. Geophys. Res. Biogeosci.* **111** 1–13
- Ping C L, Jastrow J D, Jorgenson M T, Michaelson G J and Shur Y L 2015 Permafrost soils and carbon cycling *Soil* **1** 147–71
- R Development Core Team 2015 *R: A Language and Environment for Statistical Computing* (Vienna: R Foundation for Statistical Computing)
- Rustad L E *et al* 2001 A meta-analysis of the response of soil respiration, net nitrogen mineralization, and aboveground plant growth to experimental ecosystem warming *Oecologia* **126** 543–62
- Schenk H J and Jackson R B 2002 The global biogeography of roots *Ecol. Monogr.* **72** 311–28
- Schimel J P and Bennett J 2004 Nitrogen mineralization: challenges of a changing paradigm *Ecology* **85** 591–602
- Schimel J P and Chapin F S 1996 Tundra plant uptake of amino acid and NH₄⁺ nitrogen *in situ*: plants compete well for amino acid N *Ecology* **77** 2142–7
- Schuur E A G, Crummer K G, Vogel J G and MacK M C 2007 Plant species composition and productivity following permafrost thaw and thermokarst in Alaskan Tundra *Ecosystems* **10** 280–92
- Shaver G R and Chapin F S 1980 Response to fertilization by various plant growth forms in an Alaskan tundra: nutrient accumulation and growth *Ecology* **61** 662–75
- Shaver G R and Chapin F S 1991 Production: biomass relationships and element cycling in contrasting arctic vegetation types *Ecol. Monogr.* **61** 1–31

- Shaver G R and Jonasson S 1999 Response of arctic ecosystems to climate change: results of long-term field experiments in Sweden and Alaska *Polar Res.* **18** 245–52
- Sistla S A, Moore J C, Simpson R T, Gough L, Shaver G R and Schimel J P 2013 Long-term warming restructures Arctic tundra without changing net soil carbon storage *Nature* **497** 615–8
- Sistla S A and Schimel J P 2013 Seasonal patterns of microbial extracellular enzyme activities in an arctic tundra soil: identifying direct and indirect effects of long-term summer warming *Soil Biol. Biochem.* **66** 119–29
- Sloan V L, Fletcher B J and Phoenix G K 2016 Contrasting synchrony in root and leaf phenology across multiple sub-Arctic plant communities *J. Ecol.* **104** 239–48
- Sorensen P and Jensen E S 1991 Sequential diffusion of ammonium and nitrate from soil extracts to a polytetrafluoroethylene trap for ^{15}N determination *Anal. Chim. Acta* **252** 201–3
- South A 2011 rworldmap: a new R package for mapping global data *R J.* **3** 35–43
- Soil Survey Staff 2010 *Keys to Soil Taxonomy* 11th edn (Washington, DC: United States Department of Agriculture-Natural Resources Conservation Service)
- Stark S and Väisänen M 2014 Insensitivity of soil microbial activity to temporal variation in soil N in subarctic tundra: evidence from responses to large migratory grazers *Ecosystems* **17** 906–17
- Treat C C, Wollheim W M, Varner R K and Bowden W B 2016 Longer thaw seasons increase nitrogen availability for leaching during fall in tundra soils *Environ. Res. Lett.* **11** 64013
- van Wijk M T, Gough L, Hobbie S E and Shaver G R 2003 Luxury consumption of soil nutrients: a possible competitive strategy in above-ground and below-ground biomass allocation and root morphology for slow-growing arctic vegetation? *J. Ecol.* **91** 664–76
- Vaughan D G *et al* 2013 Observations: cryosphere climate change 2013: the physical science basis *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed T F Stocker, D Qin, G K Plattner, M Tignor, S K Allen, J Boschung, A Nauels, Y Xia, V Bex and P M Midgley (Cambridge: Cambridge University Press)
- Wallenstein M D, McMahon S K and Schimel J P 2009 Seasonal variation in enzyme activities and temperature sensitivities in Arctic tundra soils *Glob. Change Biol.* **15** 1631–9
- Wanek W, Mooshammer M, Blöchl A, Hanreich A and Richter A 2010 Determination of gross rates of amino acid production and immobilization in decomposing leaf litter by a novel ^{15}N isotope pool dilution technique *Soil Biol. Biochem.* **42** 1293–302
- Wang P, Mommer L, van Ruijven J, Berendse F, Maximov T C and Heijmans M M P D 2016 Seasonal changes and vertical distribution of root standing biomass of graminoids and shrubs at a Siberian Tundra site *Plant Soil* **407** 55–65
- Weintraub M N and Schimel J P 2005 Seasonal protein dynamics in Alaskan arctic tundra soils *Soil Biol. Biochem.* **37** 1469–75
- Wild B *et al* 2016 Plant-derived compounds stimulate the decomposition of organic matter in arctic permafrost soils *Sci. Rep.* **6** 25607
- Wild B *et al* 2014 Input of easily available organic C and N stimulates microbial decomposition of soil organic matter in arctic permafrost soil *Soil Biol. Biochem.* **75** 143–51
- Wild B *et al* 2013 Nitrogen dynamics in turbic cryosols from Siberia and Greenland *Soil Biol. Biochem.* **67** 85–93
- Wild B *et al* 2015 Microbial nitrogen dynamics in organic and mineral soil horizons along a latitudinal transect in western Siberia *Glob. Biogeochem. Cycles* **29** 567–82
- Xu C, Guo L, Ping C L and White D M 2009 Chemical and isotopic characterization of size-fractionated organic matter from cryoturbated tundra soils, northern Alaska *J. Geophys. Res. Biogeosci.* **114** 1–11
- Zhu B, Gutknecht J L M, Herman D J, Keck D C, Firestone M K and Cheng W 2014 Rhizosphere priming effects on soil carbon and nitrogen mineralization *Soil Biol. Biochem.* **76** 183–92
- Zhu Q, Iversen C M, Riley W J, Slette I J and Stel H M 2016 Root traits explain observed tundra vegetation nitrogen uptake patterns: implications for trait-based land models *J. Geophys. Res. Biogeosciences* **121** 3101–12
- Zubrzycki S, Kutzbach L, Grosse G and Desyatkin A 2013 Organic carbon and total nitrogen stocks in soils of the Lena River Delta *Biogeosciences* **10** 3507–24