

# Effective and ecological half-lives of $^{90}\text{Sr}$ and $^{137}\text{Cs}$ observed in wheat and rice in Japan

Stefan Merz<sup>1</sup> · Katsumi Shozugawa<sup>2</sup> · Georg Steinhauser<sup>3,4,5</sup>

Received: 3 June 2015 / Published online: 9 August 2015  
© The Author(s) 2015. This article is published with open access at Springerlink.com

**Abstract** Published pre-Fukushima food monitoring data from 1963 to 1995 were used to study the long-term presence of  $^{137}\text{Cs}$  and  $^{90}\text{Sr}$  in rice and wheat. Effective half-lives ( $T_{\text{eff}}$ ) were calculated for rice ( $^{137}\text{Cs}$ : 5.6 years;  $^{90}\text{Sr}$ : 6.7 years) and wheat ( $^{137}\text{Cs}$ : 3.5 years;  $^{90}\text{Sr}$ : 6.2 years), respectively. In rice,  $^{137}\text{Cs}$  exhibits a longer  $T_{\text{eff}}$  because putrefaction processes will lead to the formation of  $\text{NH}_4^+$  ions that are efficient ion exchangers for mineral-adsorbed cesium ions, hence making it more readily available to the plant. Knowledge on the long-term behavior of radiocesium and radiostrontium will be important for Japanese food-safety campaigns after the Fukushima nuclear accident.

**Keywords** Fukushima · Foodstuff · Food safety · Effective half-life · Ecological half-life ·  $^{90}\text{Sr}$  ·  $^{137}\text{Cs}$

## Introduction

Nuclear accidents cause an array of health problems, both physical and mental. Fear-related problems are among the most underestimated consequences after a severe nuclear accident such as Chernobyl or Fukushima [1–3]. The lack of a sensory organ is one major reason for doubts about the own safety [4], especially when consuming potentially contaminated food. Indeed, the intake of contaminated food is the most severe pathway for exposure after nuclear accidents [5–8], which also potentially affects remote communities that have not been affected by direct fallout from the accident, but may consume contaminated food that is transported from contaminated regions.

In the aftermath of the Fukushima nuclear accident, the Japanese authorities have put much effort into a food monitoring program that stands unprecedented in human history [9]. More than 900,000 samples have been measured to date. This campaign especially also focused on rice that is traditionally a very important food item in Japanese cuisine [10]. These efforts have been supported by whole body counting campaigns of children and adult residents [11] as well as food duplicate studies [12, 13].

From a radioecological point of view, the behavior of radionuclides in the environment and the interaction (uptake, excretion) with food organisms (plants, fungi, animals) is essential for the prediction of future internal exposure due to intake of contaminated foods. Fungi, for example, have long been known to hyper-accumulate radiocesium from soil, leading to remarkable activities in certain mushroom species (and animals that feed on them). The presence and availability to the plant of a radionuclide depends on its physical half-life and its ecological half-life that is determined by wash-out and immobilization of the nuclide. The objective of this study was to determine the

---

✉ Georg Steinhauser  
georg.steinhauser@ati.ac.at;  
georg.steinhauser@colostate.edu;  
steinhauser@irs.uni-hannover.de

<sup>1</sup> Vienna University of Technology, Atominstytut,  
Stadionallee 2, 1020 Vienna, Austria

<sup>2</sup> Graduate School of Arts and Sciences, The University of  
Tokyo, Meguro-ku, Tokyo, Japan

<sup>3</sup> Environmental and Radiological Health Sciences, Colorado  
State University, Fort Collins, CO 80523, USA

<sup>4</sup> Institute of Environmental Radioactivity, Fukushima  
University, Fukushima 960-1296, Japan

<sup>5</sup> Leibniz Universität Hannover, Institut für Radioökologie und  
Strahlenschutz, 30419 Hannover, Germany

effective and ecological half-lives of  $^{137}\text{Cs}$  and  $^{90}\text{Sr}$  in both rice and wheat in Japan, as taken up from soil.

The effective half-life ( $T_{\text{eff}}$ ) combines both the physical decay as well as ecological “losses” such as migration or immobilization by adsorption on soil minerals. The ecological losses are expressed by the ecological half-life ( $T_{\text{eco}}$ ). The correlation between physical half-life ( $T_{1/2}$ ),  $T_{\text{eff}}$ , and  $T_{\text{eco}}$  is shown in Eq. 1 [14].

$$\frac{1}{T_{\text{eff}}} = \frac{1}{T_{\text{eco}}} + \frac{1}{T_{1/2}} \quad (1)$$

## Materials and methods

This study is based on pre-Fukushima (1959–1995)  $^{90}\text{Sr}$  and  $^{137}\text{Cs}$  data for polished rice [15] and polished wheat [16] that were published by Komamura et al. who studied the effect of the 20th century nuclear weapons fallout. This unique data set allows for the study of the long-term behavior of both radionuclides. What makes both papers especially valuable is that they report on the activity concentrations in rice and wheat in *Japan* (thus ideally representing the environmental setting that will be relevant for post-Fukushima challenges) and in *multiple samples* per year (thus representing a credible average). Due to its volcanic history, Japanese soil (acidic soil) is much distinct from non-volcanic soil and exhibits a unique soil chemistry. Like clay minerals, high-silica volcanic ash exhibits cation exchange properties [17].

The focus of the present study was on both radionuclides  $^{137}\text{Cs}$  ( $T_{1/2} = 30.2$  years) as well as the understudied radionuclide  $^{90}\text{Sr}$  ( $T_{1/2} = 28.6$  years) [18, 19]. We analyzed the data of nationwide averages per year published by Komamura et al. [15, 16].


## Results and discussion

The source of  $^{137}\text{Cs}$  and  $^{90}\text{Sr}$  in food (after uptake by plants from soil) has been dominated for a long time by the fallout of atmospheric nuclear explosions. Nuclear accidents, however, in particular Chernobyl and Fukushima have the potential to massively increase the fallout-background locally and regionally. In Austria, for example, the  $^{137}\text{Cs}$  inventory in soil is approximately 10 % due to weapons fallout and 90 % due to the Chernobyl accident [20, 21]. Continuous input from ongoing fallout caused by the cold war’s weapons race in the late 1950s and early 1960s led to a constantly increasing environmental inventory of  $^{90}\text{Sr}$  and  $^{137}\text{Cs}$ . A constantly increasing contamination of soil does not allow for the establishment of a certain, stable level which makes environmental processes establishing


the ecological half-life observable. Therefore, we decided to neglect data prior to 1963. In 1963, the Partial Nuclear Test-Ban Treaty was signed that prohibited atmospheric nuclear tests. In this year, the activity levels in the data base also reached a maximum for both  $^{90}\text{Sr}$  and  $^{137}\text{Cs}$ . It appeared reasonable to calculate the  $T_{\text{eff}}$  and  $T_{\text{eco}}$  from this maximum value, similar to our previous investigation of the apparent half-life of  $^{131}\text{I}$  in animal thyroids [22].

The results of the calculations are illustrated in Fig. 1 (rice) and Fig. 2 (wheat). For the exponential trend line, only data from 1963 were used for the above reason. In the data set for  $^{137}\text{Cs}$  in wheat (interestingly not in rice), an obvious outlier was observable in 1986—the year of the Chernobyl accident. Rejecting this data point greatly improved the coefficient of determination ( $R^2$ ) from 0.77 to 0.90. Rejecting the 1986  $^{90}\text{Sr}$  data point in rice did not have any effect on the  $R^2$ . This is in line with many previous observations showing that the Chernobyl nuclear accident has been a much more powerful source of  $^{137}\text{Cs}$  than of  $^{90}\text{Sr}$  [1]. Even for  $^{137}\text{Cs}$ , the following years after Chernobyl proved to follow the pre-Chernobyl pattern and hence were included in the calculation. It appears likely that, due to the long distance and maximum  $^{137}\text{Cs}$  activity concentrations in air of “only”  $0.074 \text{ Bq/m}^3$  in Japan [23], the Chernobyl nuclear accident did not significantly increase the radiocesium inventory in soil in Japan. Instead, it can be speculated that foliar uptake by the wheat plant and direct deposition of  $^{137}\text{Cs}$  particulates on the wheat grains has increased the  $^{137}\text{Cs}$  activity concentration in wheat in 1986 only.

The effective half-lives were derived from the decay constants in Figs. 1 and 2, respectively. For rice, the effective half-lives were  $T_{\text{eff}}(^{137}\text{Cs}) = 5.6$  years and  $T_{\text{eff}}(^{90}\text{Sr}) = 6.7$  years, respectively. In wheat, the effective



**Fig. 1** Activity concentrations of  $^{90}\text{Sr}$  and  $^{137}\text{Cs}$  in rice and their development over time (1959–1995) [15]. Only data from 1963 onwards were used for calculation of the exponential trend line



**Fig. 2** Activity concentrations of  $^{90}\text{Sr}$  and  $^{137}\text{Cs}$  in wheat and their development over time (1959–1995) [16]. Only data from 1963 onwards were used for calculation of the exponential trend line. For  $^{137}\text{Cs}$ , the data point from 1986 (Chernobyl) was also rejected as it represents an obvious outlier in the data set

half-lives were  $T_{\text{eff}}(^{137}\text{Cs}) = 3.5$  years and  $T_{\text{eff}}(^{90}\text{Sr}) = 6.2$  years, respectively.

From Eq. 1, the ecological half-lives were derived. In rice, they were  $T_{\text{eco}}(^{137}\text{Cs}) = 6.8$  years and  $T_{\text{eco}}(^{90}\text{Sr}) = 8.7$  years, respectively. In wheat, the ecological half-lives were  $T_{\text{eco}}(^{137}\text{Cs}) = 3.9$  years and  $T_{\text{eco}}(^{90}\text{Sr}) = 8.0$  years, respectively.

While the  $T_{\text{eff}}$  of  $^{90}\text{Sr}$  are comparable for both wheat and rice, it is interesting to note that the  $T_{\text{eff}}$  of  $^{137}\text{Cs}$  is significantly shorter for wheat than for rice. The main reason for this discrepancy is most probably the agricultural production technique in rice paddies. Putrefaction processes under water cause the formation of ammonia ( $\text{NH}_3$ ) that readily dissolves in water, thus forming ammonium ( $\text{NH}_4^+$ ) ions. These ammonium ions have been identified previously [24] as powerful ion exchangers for  $\text{Cs}^+$  ions adsorbed on (clay) minerals. Other reasons for explaining this discrepancy may be different plant-related uptake mechanisms and kinetics for  $\text{Sr}^{2+}$  and  $\text{Cs}^+$ , respectively and different translocation inside the plant.

After the Fukushima nuclear accident, the authorities assumed an intrinsic coexistence of  $^{137}\text{Cs}$  and  $^{90}\text{Sr}$  and thus a constant  $^{90}\text{Sr}/^{137}\text{Cs}$  ratio in foods. The maximum contribution of  $^{90}\text{Sr}$  was assumed to be  $\leq 10\%$  (after April 2012  $\leq 0.3\%$ ) of the respective  $^{137}\text{Cs}$  activity concentration in food. This allowed for the rapid determination of  $^{137}\text{Cs}$  using gamma-spectrometry and the subsequent estimation of the dose contribution caused by  $^{90}\text{Sr}$  (a radionuclide that requires a very laborious and time-consuming separation prior to radiometric analysis). In our previous study [9] we discovered that the assumption of a constant  $^{90}\text{Sr}/^{137}\text{Cs}$  ratio is not justified over longer periods

of time. In agreement with our previous conclusions we can now specify that the different effective ecological half-lives of both radionuclides thwart the assumption of a constant ratio over time, thus potentially putting the reliability and public credibility of the food monitoring program at stake.

## Conclusions

We presented evidence for significantly different effective half-lives of  $^{137}\text{Cs}$  (3.5 years) and  $^{90}\text{Sr}$  (6.2 years) in wheat, which will be of great importance for food safety considerations. In rice, in contrast, the effective half-lives of  $^{137}\text{Cs}$  (5.6 years) and  $^{90}\text{Sr}$  (6.7 years) proved to be rather comparable, which can probably be explained by exceptional properties of the agricultural production technique in a paddy. This technique, in particular, the formation of  $\text{NH}_4^+$  ions (superior ion exchangers), makes radiocesium more bioavailable to the rice plant compared to the wheat plant. The longer  $T_{\text{eff}}$  of  $^{137}\text{Cs}$  observed in rice hence should be considered as an ecological exception and should not be applied on other food items without experimental evidence. The ecological half-lives in rice were  $T_{\text{eco}}(^{137}\text{Cs}) = 6.8$  years and  $T_{\text{eco}}(^{90}\text{Sr}) = 8.7$  years, respectively. For wheat,  $T_{\text{eco}}(^{137}\text{Cs}) = 3.9$  years and  $T_{\text{eco}}(^{90}\text{Sr}) = 8.0$  years were calculated. In both rice and wheat, the ecological half-lives of  $^{90}\text{Sr}$  were longer compared to  $^{137}\text{Cs}$ , which is most likely due to the higher bioavailability of strontium in soil, because cesium ions have a higher affinity to clay minerals and become immobilized more readily.

**Acknowledgments** The authors are indebted to M. Komamura and colleagues whose pioneering work has made this study possible. This work was supported by JSPS KAKENHI (Grant Number 25870158), from CDC NIOSH Mountain and Plains Education and Research Center (Grant Number T42OH009229-07), and the US Nuclear Regulatory Commission (NRC) (Grant Number NRC-HQ-12-G-38-0044). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the CDC NIOSH and MAP ERC.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Steinhauser G, Brandl A, Johnson TE (2014) Comparison of the Chernobyl and Fukushima nuclear accidents: a review of the environmental impacts. *Sci Total Environ* 470–471:800–817

2. Shigemura J, Tanigawa T, Saito I, Nomura S (2012) Psychological distress in workers at the Fukushima nuclear power plants. *J Am Med Assoc* 308:667–669
3. Bromet EJ (2012) Mental health consequences of the Chernobyl disaster. *J Radiol Prot* 32:N71–N75
4. Steinhauser G (2015) Ionizing radiation—an evolutionary threat? *Hypothesis* 13(1):e6. doi:10.5779/hypothesis.v13i1.437
5. Travnikova IG, Bruk GJ, Shutov VN, Bazjukin AB, Balonov MI, Rahola T, Tillander M (2001) Contribution of different foodstuffs to the internal exposure of rural inhabitants in Russia after the Chernobyl accident. *Radiat Prot Dosim* 93:331–339
6. Hamada N, Ogino H (2012) Food safety regulations: what we learned from the Fukushima nuclear accident. *J Environ Radioact* 111:83–99
7. Hamada N, Ogino H, Fujimichi Y (2012) Safety regulations of food and water implemented in the first year following the Fukushima nuclear accident. *J Radiat Res* 53:641–671
8. Merz S, Steinhauser G, Hamada N (2013) Anthropogenic radionuclides in Japanese food: environmental and legal implications. *Environ Sci Technol* 47:1248–1256
9. Merz S, Shozugawa K, Steinhauser G (2015) Analysis of Japanese radionuclide monitoring data of food before and after the Fukushima nuclear accident. *Environ Sci Technol* 49:2875–2885
10. Nihei N, Tanoi K, Nakanishi TM (2015) Inspections of radiocesium concentration levels in rice from Fukushima Prefecture after the Fukushima Dai-ichi Nuclear Power Plant accident. *Sci Rep* 5:8653
11. Hayano RS, Tsubokura M, Miyazaki M, Satou H, Sato K, Sakuma Y (2013) Internal radiocesium contamination of adults and children in Fukushima 7 to 20 months after the Fukushima NPP accident as measured by extensive whole-body-counter survey. *Proc Jpn Acad Ser B* 89:157–163
12. Harada KH, Fujii Y, Adachi A, Tsukidate A, Asai F, Koizumi A (2012) Dietary intake of radiocesium in adult residents in Fukushima prefecture and neighboring regions after the Fukushima nuclear power plant accident: 24-h food-duplicate survey in December 2011. *Environ Sci Technol* 47:2520–2526
13. Koizumi A, Harada KH, Niisoe T, Adachi A, Fujii Y, Hitomi T, Kobayashi H, Wada Y, Watanabe T, Ishikawa H (2012) Preliminary assessment of ecological exposure of adult residents in Fukushima Prefecture to radioactive cesium through ingestion and inhalation. *Environ Health Prev Med* 17:292–298
14. Zibold G, Klemm E (2005) Ecological half-times of  $^{137}\text{Cs}$  and  $^{90}\text{Sr}$  in forest and freshwater ecosystems. *Radioprotection* 40:S497–S502
15. Komamura M, Tsumura A, Kodaira K (2001)  $^{90}\text{Sr}$  and  $^{137}\text{Cs}$  contamination of polished rice in Japan. Survey and analysis during the years 1959–1995. *Radioisotopes* 50:80–93 (**in Japanese**)
16. Komamura M, Tsumura A, Kihou N, Kodaira K (2002)  $^{90}\text{Sr}$  and  $^{137}\text{Cs}$  contamination of wheat produced in Japan. Survey and analysis during the years 1959 through 1995 including the Chernobyl accident. *Radioisotopes* 51:345–363 (**in Japanese**)
17. Steinhauser G, Bichler M (2008) Adsorption of ions onto high-silica volcanic glass. *App Radiat Isot* 66:1–8
18. Steinhauser G (2014) Fukushima's forgotten radionuclides: a review of the understudied radioactive emissions. *Environ Sci Technol* 48:4649–4663
19. Steinhauser G, Schauer V, Shozugawa K (2013) Concentration of strontium-90 at selected hot spots in Japan. *PLoS One* 8:e57760
20. Steinhauser G, Merz S, Hainz D, Sterba JH (2013) Artificial radioactivity in environmental media (air, rainwater, soil, vegetation) in Austria after the Fukushima nuclear accident. *Environ Sci Pollut Res* 20:2527–2537
21. Bossew P, Falkner T, Henrich E, Kienzl K (1995) Cäsiumbelastung der Böden Österreichs (Contamination of Austrian soils by Caesium-137). Umweltbundesamt, Wien (**in German**)
22. Steinhauser G, Merz S, Kübber-Heiss A, Katzlberger C (2012) Using animal thyroids as ultra-sensitive biomonitors for environmental radioiodine. *Environ Sci Technol* 46:12890–12894
23. Higuchi H, Fukatsu H, Hashimoto T, Nonaka N, Yoshimizu K, Omine M et al (1988) Radioactivity in surface air and precipitation in Japan after the Chernobyl accident. *J Environ Radioact* 6:131–144
24. Kodaira K (1964) Radioactive contamination of rice in Japan with reference to Sr-90 and Cs-137 content in rice until 1962. *J Radiat Res* 39:116–119