

ISOGENIES OF PRYM VARIETIES

ROBERTO LAFACE - CÉSAR MARTÍNEZ

We prove an extension of the Babbage-Enriques-Petri theorem for semi-canonical curves. We apply this to show that the Prym variety of a generic element of a codimension k subvariety of \mathcal{R}_g is not isogenous to another distinct Prym variety, under some mild assumption on k .

1. Introduction

Let \mathcal{R}_g denote the moduli space of unramified irreducible double covers of complex smooth curves of genus g . Given an element $\pi : D \rightarrow C$ in \mathcal{R}_g , we can lift this morphism to the corresponding Jacobians via the norm map

$$\mathrm{Nm}_\pi : J(D) \rightarrow J(C).$$

By taking the neutral connected component of its kernel, we obtain an abelian variety of dimension $g - 1$ called the *Prym variety* attached to π .

In this note, we study the isogeny locus in \mathcal{A}_{g-1} of Prym varieties attached to generic elements in \mathcal{R}_g ; that is, principally polarized abelian varieties of dimension $g - 1$ which are isogenous to such Prym varieties. More concretely, given a subvariety \mathcal{Z} of \mathcal{R}_g of codimension k and a generic element $\pi : D \rightarrow C$ in \mathcal{Z} , we prove that the Prym variety attached to π is not isogenous to a distinct Prym variety, whenever $g \geq \max\{7, 3k + 5\}$, see Theorem 3.2.

Entrato in redazione: 31 maggio 2017

AMS 2010 Subject Classification: C0D1C3, PR0V4

Keywords: abelian varieties, Prym varieties, isogenies, semi-canonical curves

This result can be seen as an extension of the analogue statements for Jacobians of generic curves proven by Bardelli and Pirola [1] for the case $k = 0$, and Marcucci, Naranjo and Pirola [2] for $k > 0$, $g \geq 3k + 5$ or $k = 1$ and $g \geq 5$. In the latter, to prove the case $g \geq 3k + 5$, they use an argument on infinitesimal variation of Hodge structure proposed by Voisin in [1, Remark (4.2.5)] which allows them to translate the question to a geometric problem of intersection of quadrics. In doing so, they give a generalization of Babbage-Enriques-Petri's theorem which allows them to recover a canonical curve from the intersection of a system of quadrics in \mathbb{P}^{g-1} of codimension k . The strategy we follow to prove Theorem 3.2 is an adaptation of these techniques to the setting of Prym varieties. We are also able to give an extension of Babbage-Enriques-Petri's theorem for semicanonical curves in a similar fashion as in [2], see Proposition 2.2. Our result generalises the one by Lange and Sernesi [3] for curves of genus $g \geq 9$, since it recovers a semicanonical curve of genus $g \geq 7$ from a system of quadrics in \mathbb{P}^{g-2} of codimension k , $g \geq 3k + 5$.

2. Intersection of quadrics

Let C be a smooth curve. Given a globally generated line bundle $L \in \text{Pic}(C)$, we denote by $\varphi_L : C \rightarrow \mathbb{P}H^0(C, L)^*$ its induced morphism. If L is very ample, we say that $\varphi_L(C)$ is *projectively normal* if its homogeneous coordinate ring is integrally closed; or equivalently, if for all $k \geq 0$, the homomorphism

$$\text{Sym}^k H^0(L) \longrightarrow H^0(L^{\otimes k})$$

is surjective.

We also recall that the *Clifford index* of C is defined as

$$\min\{\deg(L) - 2h^0(C, L) + 2\},$$

where the minimum ranges over the line bundles $L \in \text{Pic}(C)$ such that $h^0(C, L) \geq 2$ and $h^0(C, \omega_C \otimes L^{-1}) \geq 2$. Its value is an integer between 0 and $\lfloor \frac{g-1}{2} \rfloor$, where g is the genus of the curve.

Let C be of genus g and with Clifford index c . For any non-trivial 2-torsion point η in the Jacobian of C , we call $\omega_C \otimes \eta$ a *semicanonical line bundle* of C whenever it is globally generated, and we denote by $\varphi_{\omega_C \otimes \eta} : C \rightarrow \mathbb{P}^{g-2}$ its associated morphism. In that case, we call its image $C_\eta := \varphi_{\omega_C \otimes \eta}(C)$ a *semicanonical curve*. The following is a result of Lange and Sernesi [3], and Lazarsfeld [4]:

Lemma 2.1. *If $g \geq 7$ and $c \geq 3$, then $\omega_C \otimes \eta$ is very ample and the semicanonical curve C_η is projectively normal.*

Furthermore, Lange and Sernesi prove that C_η is the only non-degenerate curve in the intersection of all quadrics in \mathbb{P}^{g-2} containing C_η if $c > 3$, or $c = 3$ and $g \geq 9$, see [3]. The following proposition generalises this result for a smaller family of quadrics.

Proposition 2.2. *Let C be a curve of genus g and Clifford index c , and η be a non-trivial 2-torsion point in $J(C)$. Let $I_2(C_\eta) \subset \text{Sym}^2 H^0(C, \omega_C \otimes \eta)$ be the vector space of equations of the quadrics containing C , and $K \subset I_2(C_\eta)$ be a linear subspace of codimension k . If $g \geq \max\{7, 2k+6\}$ and $c \geq \max\{3, k+2\}$, then C_η is the only irreducible non-degenerate curve in the intersection of the quadrics of K .*

Notice that for $k = 0$, this proposition extends the result of Lange and Sernesi [3] to the cases when $c = 3$ and $g = 7$ and 8 . We refer to Remark 2.3 for a brief discussion on a simplified version of the following proof in this case.

Proof. We start by assuming that there exists an irreducible non-degenerate curve C_0 in the intersection of quadrics $\bigcap_{Q \in K} Q \subset \mathbb{P}H^0(C, \omega_C \otimes \eta)^*$, which is different from C_η . In particular, we can choose $k+1$ linearly independent points in $\bigcap_{Q \in K} Q$ such that $x_i \notin C_\eta$ for all i . By abuse of notation, we denote also as x_i the representatives in $H^0(C, \omega_C \otimes \eta)^*$. We define $L \subset \text{Sym}^2 H^0(C, \omega_C \otimes \eta)^*$ as the linear subspace spanned by $x_i \otimes x_i$.

Let $R = I_2(C_\eta)/K$ and $R' = \text{Sym}^2 H^0(C, \omega_C \otimes \eta)/K$. By Lemma 2.1 and the fact that $g \geq 7$ and $c \geq 3$, we have that C_η is projectively normal. Hence, we can build the following diagram:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & K & \longrightarrow & I_2(C_\eta) & \longrightarrow & R \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \\
 & & \text{Sym}^2 H^0(C, \omega_C \otimes \eta) = \text{Sym}^2 H^0(C, \omega_C \otimes \eta) & & & & \\
 & & \downarrow & & \downarrow & & \\
 0 \longrightarrow R & \longrightarrow & R' & \longrightarrow & H^0(C, \omega_C^{\otimes 2}) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & &
 \end{array}$$

where the last row is obtained by applying the snake lemma to the first two rows.

By dualizing this diagram, we get

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & \mathrm{H}^0(C, \omega_C^{\otimes 2})^* = \mathrm{H}^1(C, T_C) & \longrightarrow & R'^* & \longrightarrow & R^* \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \\
 & & \mathrm{Sym}^2 \mathrm{H}^0(C, \omega_C \otimes \eta)^* = \mathrm{Sym}^2 \mathrm{H}^0(C, \omega_C \otimes \eta)^* & & & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & R^* & \longrightarrow & I_2(C_\eta)^* & \longrightarrow & K^* \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & &
 \end{array}$$

Notice that $Q(\alpha) = 0$ for every $\alpha \in L$ and every $Q \in K$. Therefore, $L \subset R'^*$. Since $\dim(L) = k + 1$ and $\dim(R) = k$, there is a non-trivial element $\alpha \in L \cap \mathrm{H}^1(C, T_C)$. By the isomorphism $\mathrm{H}^1(C, T_C) \simeq \mathrm{Ext}^1(\omega_C, \mathcal{O}_C)$, there is a 2 vector bundle E_α associated to α satisfying the following exact sequence:

$$0 \longrightarrow \mathcal{O}_C \longrightarrow E_\alpha \longrightarrow \omega_C \longrightarrow 0. \quad (1)$$

The cup product with α is the coboundary map $\mathrm{H}^0(C, \omega_C) \rightarrow \mathrm{H}^1(C, \mathcal{O}_C)$. By writing the element $\alpha = \sum_{i=1}^{k+1} a_i x_i \otimes x_i$, we have

$$\mathrm{Ker}(\cdot \cup \alpha) = \bigcap_{i \mid a_i \neq 0} H_i,$$

where $H_i = \mathrm{Ker}(x_i)$. After reordering, we may assume that $x_1, \dots, x_{k'}$ are the points such that $a_i \neq 0$, for some $k' \leq k + 1$. This means that there are $g - k'$ linearly independent sections in $\mathrm{H}^0(C, \omega_C)$ lifting to $\mathrm{H}^0(C, E_\alpha)$. Denote by $W \subset \mathrm{H}^0(C, E_\alpha)$ the vector space generated by these sections, and consider the morphism $\psi: \wedge^2 W \rightarrow \mathrm{H}^0(C, \omega_C)$ obtained by the following composition:

$$\wedge^2 W \longrightarrow \wedge^2 \mathrm{H}^0(C, E_\alpha) \longrightarrow \mathrm{H}^0(C, \det E_\alpha) = \mathrm{H}^0(C, \omega_C).$$

The kernel of ψ has codimension at most g , and the Grassmannian of the decomposable elements in $\mathbb{P}(\wedge^2 W)$ has dimension $2(g - k' - 2)$. Since $g > 2k + 5$ by hypothesis, we have that their intersection is not trivial. Thus, take $s_1, s_2 \in \mathrm{H}^0(C, E_\alpha)$ such that $\psi(s_1 \wedge s_2) = 0$. They generate a line bundle $M_\alpha \subset E_\alpha$ and $h^0(C, M_\alpha) \geq 2$. Take Q_α the neutral component of the quotient E_α/M_α , and L_α the kernel of $E_\alpha \rightarrow Q_\alpha$, then we obtain the following exact sequence:

$$0 \longrightarrow L_\alpha \longrightarrow E_\alpha \longrightarrow Q_\alpha \longrightarrow 0. \quad (2)$$

Notice that $M_\alpha \subset L_\alpha$, hence $h^0(C, L_\alpha) \geq 2$. Moreover from (1) and (2), we obtain $\omega_C \simeq \det E_\alpha \simeq L_\alpha \otimes Q_\alpha$, which implies that $Q_\alpha \simeq \omega_C \otimes L_\alpha^{-1}$. We have the following diagram:

$$\begin{array}{ccccccc}
 & & & 0 & & & \\
 & & & \downarrow & & & \\
 & & & L_\alpha & & & \\
 & & & \downarrow & & & \\
 0 & \longrightarrow & \mathcal{O}_C & \longrightarrow & E_\alpha & \longrightarrow & \omega_C \longrightarrow 0 \\
 & & \searrow & & \downarrow & & \\
 & & & & \omega_C \otimes L_\alpha^{-1} & & \\
 & & & & \downarrow & & \\
 & & & & 0 & &
 \end{array}$$

Assume that $\mathcal{O}_C \rightarrow \omega_C \otimes L_\alpha^{-1}$ is 0. Then the section of E_α that represents $\mathcal{O}_C \rightarrow E_\alpha$ would be a section of L_α , in particular, a section in W . Since the sections in W map to sections of ω_C , this contradicts the exactness of the horizontal sequence. So $\mathcal{O}_C \rightarrow \omega_C \otimes L_\alpha^{-1}$ is not 0 and the $h^0(C, \omega_C \otimes L_\alpha^{-1}) > 0$.

If $h^0(C, \omega_C \otimes L_\alpha^{-1}) \geq 2$, we have that

$$c \leq \deg(L_\alpha) - 2h^0(C, L_\alpha) + 2. \tag{3}$$

Moreover, $h^0(C, L_\alpha) + h^0(C, \omega_C \otimes L_\alpha^{-1}) \geq h^0(C, E_\alpha) > \dim(W) = g - k'$ and, using Riemann-Roch we obtain that $2h^0(C, L_\alpha) \geq \deg(L_\alpha) + 2 - k'$. Combining this with (3), we obtain that $c \leq k' \leq k + 1$ which contradicts our hypothesis on c ($c \geq k + 2$). Hence, $h^0(C, \omega_C \otimes L_\alpha^{-1}) = 1$.

Write $\omega_C \otimes L_\alpha^{-1} \simeq \mathcal{O}_C(p_1 + \dots + p_e)$, where $e = \deg(\omega_C \otimes L_\alpha^{-1})$. Notice that $h^0(C, L_\alpha) \geq g - k'$ and $\deg(L_\alpha) = 2g - 2 - e$. Using Riemann-Roch, we get

$$g - k' \leq h^0(C, L_\alpha) = h^0(C, \omega_C \otimes L_\alpha^{-1}) + 2g - 2 - e - (g - 1) = g - e.$$

So $e \leq k'$.

By (2), we have that $L_\alpha \simeq \omega_C(-p_1 - \dots - p_e)$. Moreover, the sections of L_α lie in W , and by construction of W we have that $H^0(\omega_C(-p_1 - \dots - p_e)) \subset \text{Ker}(\cdot \cup \alpha) = \cap_{i|a_i \neq 0} H_i$. Therefore, by dualizing this inclusion, we obtain that

$$\langle x_1, \dots, x_{k'} \rangle_{\mathbb{C}} \subset \langle p_1, \dots, p_e \rangle_{\mathbb{C}}. \tag{4}$$

Let $\gamma : N_0 \rightarrow C_0$ be a normalization. For any generic choice of $k + 1$ points $x_i \in N_0$, we can repeat the construction above for $\gamma(x_1), \dots, \gamma(x_{k+1})$, and we can

assume that k' and e are constant. We define the correspondence

$$\Gamma = \left\{ (x_1 + \dots + x_{k'}, p_1 + \dots + p_e) \in N_0^{(k')} \times C_\eta^{(e)}, \right. \\ \left. \text{such that } \langle \gamma(x_1), \dots, \gamma(x_{k'}) \rangle_{\mathbb{C}} \subset \langle p_1, \dots, p_e \rangle_{\mathbb{C}} \right\}.$$

Observe that Γ dominates $N_0^{(k')}$, so $e \leq k' \leq \dim \Gamma$. In addition, the second projection $\Gamma \rightarrow C_\eta^{(e)}$ has finite fibers, since both curves are non-degenerate. This implies that $\dim \Gamma \leq e$, and so we have $k' = e$. Since $k' \leq k + 1 \leq g - 3$, by the uniform position theorem we have that the rational maps

$$C^{(k')} \dashrightarrow \text{Sec}^{(k')}(C_\eta) \subset \mathbb{G}(e - 1, \mathbb{P}^{g-2}), \\ N_0^{(k')} \dashrightarrow \text{Sec}^{(k')}(N_0) \subset \mathbb{G}(e - 1, \mathbb{P}^{g-2}),$$

are generically injective. This gives a birational map between $C_\eta^{(k')}$ and $N_0^{(k')}$. In particular, it induces dominant morphisms $J C_\eta \rightarrow J N_0$ and $J N_0 \rightarrow J C_\eta$. Therefore, $g(C_\eta) = g(N_0)$ and by a theorem of Ran [5], the birational map $C_\eta^{(k')} \dashrightarrow N_0^{(k')}$ is defined by a birational map between C_η and N_0 . By composing it with the normalization map γ , we obtain a birational map

$$\varphi : C_\eta \dashrightarrow C_0,$$

that defines the correspondence Γ ; that is $\langle \varphi(x_1), \dots, \varphi(x_{k'}) \rangle = \langle x_1, \dots, x_{k'} \rangle$ for generic elements $x_1 + \dots + x_{k'} \in C_\eta^{(k')}$. This implies that φ is generically the identity map over C_η . Thus $C_\eta = C_0$, which is a contradiction and ends the proof. \square

Remark 2.3. The proof of Corollary 3.1 can be simplified for the case $K = I_2(C_\eta)$, that is $k = 0$. Under this assumption, we only consider one point $x \notin C_\eta$, and $k' = e = 1$. Therefore, the inclusion (4) already implies the equality $C_\eta = C_0$.

3. Main theorem

An element in \mathcal{R}_g can be identified with a pair (C, η) , where C is a complex smooth curve of genus g , and η is a non-trivial 2-torsion element in the Jacobian of C . This allows us to consider \mathcal{R}_g as a covering of the moduli space \mathcal{M}_g of complex smooth curves of genus g . It is given by the morphism

$$\mathcal{R}_g \longrightarrow \mathcal{M}_g, \quad (C, \eta) \longmapsto C,$$

which has degree $2^{2g} - 1$. Thus, a generic choice of an element in a subvariety $\mathcal{Z} \subset \mathcal{R}_g$ is equivalent to a generic choice of a curve C in the image of \mathcal{Z} in \mathcal{M}_g , and any non-trivial element $\eta \in J(C)[2]$.

The following result is a direct consequence of Proposition 2.2 and it is the version of Babbage-Enriques-Petri's theorem that we use in the proof of the main result in this article.

Corollary 3.1. *Let (C, η) be a generic point in a subvariety \mathcal{Z} of \mathcal{R}_g of codimension k . Let $I_2(C_\eta) \subset \text{Sym}^2 H^0(C, \omega_C \otimes \eta)$ be the vector space of the equations of quadrics in \mathbb{P}^{g-2} containing C_η . Let $K \subset I_2(C_\eta)$ be a linear subspace of codimension k . If $g \geq \max\{7, 3k + 5\}$, then C_η is the only irreducible non-degenerate curve in the intersection of the quadrics of K .*

Proof. Let \mathcal{M}_g^c be the locus in \mathcal{M}_g corresponding to curves with Clifford index c . Then \mathcal{M}_g^c is a finite union of subvarieties of \mathcal{M}_g , where the one of higher dimension corresponds to the curves whose Clifford index is realized by a g_{c+2}^1 linear series, see [7]. By Riemann-Hurwitz, the codimension in \mathcal{M}_g of the component of the curves with a g_{c+2}^1 linear series is

$$3g - 3 - (2g - 2c + 2 - 3) = g - 2c - 2.$$

If $k = 0$, a generic curve in \mathcal{M}_g has Clifford index $c \geq 3$, because $g \geq 7$. As when $k > 0$, since $g \geq 3k + 5$, we obtain

$$k \geq g - 2c - 2 \geq 3k + 5 - 2c - 2 = 3k - 2c + 3,$$

and thus $c \geq k + 2$. The corollary follows by applying Proposition 2.2. □

Let $\widetilde{\mathcal{A}}_g^m$ be the space of isogenies of principally polarized Abelian varieties of degree m (up to isomorphism); that is the space of classes of isogenies $\chi : A \rightarrow A'$ such that $\chi^* L_{A'} \cong L_A^{\otimes m}$, where L_A (respectively $L_{A'}$) is a principal polarization on A (respectively A'). There are two forgetful maps to the moduli space \mathcal{A}_g of p.p.a.v. of dimension g

$$\begin{array}{ccc}
 & \widetilde{\mathcal{A}}_g^m & \\
 \varphi \swarrow & & \searrow \psi \\
 \mathcal{A}_g & & \mathcal{A}_g,
 \end{array}
 \tag{5}$$

such that $\varphi(\chi) = (A, L_A)$ and $\psi(\chi) = (A', L_{A'})$. These maps yield the following

commutative diagram,

$$\begin{array}{ccc}
 & T_{[\chi]}\widetilde{\mathcal{A}}_{g-1}^m & \\
 d\varphi \swarrow & & \searrow d\psi \\
 T_{[A]}\mathcal{A}_{g-1} & \xrightarrow{\lambda} & T_{[A']}\mathcal{A}_{g-1}
 \end{array} \tag{6}$$

where all maps are isomorphisms.

Theorem 3.2. *Let $\mathcal{Z} \subset \mathcal{R}_g$ be a (possibly reducible) codimension k subvariety. Assume that $g \geq \max\{7, 3k + 5\}$, and let (C, η) be a generic element in \mathcal{Z} . If there is a pair $(C', \eta') \in \mathcal{R}_g$ such that there exists an isogeny $\chi : P(C, \eta) \rightarrow P(C', \eta')$, then $(C, \eta) \cong (C', \eta')$ and $\chi = [n]$, for some $n \in \mathbb{Z}$.*

Proof. Suppose that (C, η) is generic in \mathcal{Z} . By the assumption on g , the Clifford index of a generic element of \mathcal{Z} is at least three (as shown in the proof of Corollary 3.1). However, by [8], if the Clifford index of a curve C is $c \geq 3$, then the corresponding fiber of the Prym map is 0-dimensional, i.e. $\dim P^{-1}(P(C, \eta)) = 0$. Therefore, the restriction of the Prym map to \mathcal{Z} ,

$$P|_{\mathcal{Z}} : \mathcal{Z} \rightarrow \mathcal{R}_g \rightarrow \mathcal{A}_{g-1},$$

has generically fixed degree d onto its image, for some $d \in \mathbb{N}$. So, by the genericity of the pair (C, η) , we can assume that (C, η) lies in the locus of \mathcal{Z} where $P|_{\mathcal{Z}}$ is étale. This gives the isomorphisms of the tangent spaces

$$T_{P[(C,\eta)]}P(\mathcal{Z}) \cong T_{[C,\eta]}\mathcal{Z} \quad \text{and} \quad T_{P[(C,\eta)]}P(\mathcal{R}_g) \cong T_{[C,\eta]}\mathcal{R}_g. \tag{7}$$

Let us assume that the locus of curves in \mathcal{R}_g whose corresponding Prym variety is isogenous to the Prym variety of an element in \mathcal{Z} has an irreducible component \mathcal{Z}' of codimension k . By [6], since $k < g - 2$, we have $\text{End}(P(C, \eta)) \cong \mathbb{Z}$. Suppose that we are given an isogeny $\chi : P(C, \eta) \rightarrow P(C', \eta')$; then, it must have the property that the pull-back of the principal polarization Ξ' is a multiple of the principal polarization Ξ on $P(C, \eta)$, say $\chi^*\Xi' \cong \Xi^{\otimes m}$, for some $m \in \mathbb{Z}$.

For such m , we have the diagram of forgetful maps as in (5) with $g - 1$ in place of g . We can find an irreducible subvariety $\mathcal{V} \subset \widetilde{\mathcal{A}}_{g-1}^m$ which dominates both $P(\mathcal{Z})$ and $P(\mathcal{Z}')$ through φ and ψ respectively. Setting $\mathcal{R} := \varphi^{-1}(P(\mathcal{R}_g))$ and $\mathcal{R}' := \psi^{-1}(P(\mathcal{R}_g))$, we have the inclusion $\mathcal{V} \subset \mathcal{R} \cap \mathcal{R}'$.

For a generic element $\chi : P(C, \eta) \rightarrow P(C', \eta')$ in \mathcal{V} , the diagram (6) becomes

$$\begin{array}{ccc}
 & T_{[\chi]}\widetilde{\mathcal{A}}_{g-1}^m & \\
 d\varphi \swarrow & & \searrow d\psi \\
 T_{[P(C,\eta)]}\mathcal{A}_{g-1} & \xrightarrow[\cong]{\lambda} & T_{[P(C',\eta')]\mathcal{A}_{g-1}}
 \end{array}$$

In addition, $T_{[P(C,\eta)]}\mathcal{A}_{g-1} \cong \text{Sym}^2 H^0(P(C, \eta), T_{P(C,\eta)}) \cong \text{Sym}^2 H^0(\omega_C \otimes \eta)^*$. By looking at $d\phi$, and the isomorphisms in (7), we see that we have the following diagram of tangents spaces and identifications:

$$\begin{array}{ccccccc}
 T_{[\chi]}\mathcal{V} & \longrightarrow & T_{[\chi]}\mathcal{R} & \longrightarrow & T_{[\chi]}\mathcal{R} + T_{[\chi]}\mathcal{R}' & \longrightarrow & T_{[\chi]}\widetilde{\mathcal{A}}_{g-1}^m \\
 \cong \downarrow & & \cong \downarrow & & \parallel & & \cong \downarrow \\
 T_{[C,\eta]}\mathcal{Z} & \longrightarrow & T_{C,\eta}\mathcal{R}_g & \longrightarrow & \bar{T} & \longrightarrow & \text{Sym}^2 H^0(\omega_C \otimes \eta)^*
 \end{array}$$

where the vertical arrows are $d\phi$.

By the Grassmann formula, $\dim \bar{T} \leq 3g - 3 + k$. Set

$$K(C_\eta) := \ker \left(\text{Sym}^2 H^0(\omega_C \otimes \eta) \longrightarrow \bar{T}^* \right).$$

It is a subspace of the space of quadrics containing the semicanonical curve C_η . Notice that $\text{codim}_{I_2(C_\eta)} K(C_\eta) \leq k$. By repeating the above argument with ψ in place of ϕ , we get the corresponding inclusion of vector spaces $K(C'_{\eta'}) \subset I_2(C'_{\eta'})$, and by using the (canonical) isomorphism λ above, we get a (canonical) isomorphism $K(C_\eta) \cong K(C'_{\eta'})$.

A closer look at $\lambda : T_{[P(C,\eta)]}\mathcal{A}_{g-1} \longrightarrow T_{[P(C',\eta')]\mathcal{A}_{g-1}}$ reveals that this map is induced by the isogeny $\chi : P(C, \eta) \longrightarrow P(C', \eta')$. In fact, one has that $d_0\chi : H^0(\omega_C \otimes \eta) \longrightarrow H^0(\omega_{C'} \otimes \eta')$ is an isomorphism, and λ is induced by it. This means that $d_0\chi$ induces an isomorphism of projective spaces $\mathbb{P}H^0(\omega_C \otimes \eta)^* \longrightarrow \mathbb{P}H^0(\omega_{C'} \otimes \eta')^*$, which sends quadrics containing $C'_{\eta'}$ to quadrics containing C_η , by means of λ . By using Lemma 3.1, we get that $C_\eta \cong C'_{\eta'}$, and thus $C \cong C'$. This gives us the following commutative diagram

$$\begin{array}{ccccc}
 C & \xrightarrow{\phi_{\omega_C \otimes \eta}} & C_\eta & \longrightarrow & \mathbb{P}H^0(\omega_C \otimes \eta)^* \\
 \cong \downarrow & & \cong \downarrow & & \cong \downarrow \\
 C' & \xrightarrow{\phi_{\omega_{C'} \otimes \eta'}} & C'_{\eta'} & \longrightarrow & \mathbb{P}H^0(\omega_{C'} \otimes \eta')^*
 \end{array}$$

from which we deduce that $(C, \eta) \cong (C', \eta')$. Indeed, pulling back hyperplanes to C and C' , yields an isomorphism $\omega_{C'} \otimes \eta' \cong \omega_C \otimes \eta$, from which it follows that $\eta \cong \eta'$. The isogeny is necessarily of the form $[n]$, for some $n \in \mathbb{Z}$, because $\text{End}(P(C, \eta)) \cong \mathbb{Z}$. □

Acknowledgements

We warmly thank the organizing committee of Pragmatic for the generous funding and the opportunity of enjoying such a lively and stimulating environment.

We thank Juan Carlos Naranjo and Víctor González Alonso for proposing this problem and for their helpful advice.

REFERENCES

- [1] F. Bardelli and G. P. Pirola, *Curves of genus g lying on a g -dimensional Jacobian variety*, *Inventiones Mathematicae* 95 (2) (1989), 263–276.
- [2] V. Marcucci, J. C. Naranjo and G. P. Pirola, *Isogenies of Jacobians*, To appear in *Algebraic Geometry*.
- [3] H. Lange and E. Sernesi, *Quadrics containing a Prym-canonical curve*, *Journal of Algebraic Geometry* 5 (2) (1996), 387–399.
- [4] R. Lazarsfeld, *A sampling of vector bundle techniques in the study of linear series*, *Lectures on Riemann surfaces* (Trieste, 1987), 500–559, World Sci. Publ., Teaneck, NJ (1989).
- [5] Z. Ran, *On a theorem of Martens*, *Rend. Sem. Mat. Univ. Politec. Torino* 44 (2) (1986), 287–291.
- [6] G. P. Pirola, *Base number theorem for abelian varieties. An infinitesimal approach*, *Math. Ann.* 282 (3) (1988), 361–368.
- [7] M. Coppens, G. Martens, *Secant spaces and Clifford's theorem*, *Compositio Math.* 78 (2) (1991), 193–212.
- [8] J. C. Naranjo, *The positive dimensional fibres of the Prym map*, *Pac. J. Math.* 172 (1) (1996), 223–226.

ROBERTO LAFACE

Institut für Algebraische Geometrie

Leibniz Universität Hannover

Welfengarten 1, 30167 Hannover (Germany)

e-mail: laface@math.uni-hannover.de

Technische Universität München

Zentrum Mathematik - M11

Boltzmannstraße 3, 85478 Garching bei München (Germany)

e-mail: laface@ma.tum.de

CÉSAR MARTÍNEZ

Laboratoire de Mathématiques Nicolas Oresme

Université de Caen Normandie BP 5186

14032 Caen Cedex (France)

e-mail: cesar.martinez@unicaen.fr